

Probability Theory 2 (NMSA405)

Zbyněk Pawlas

December 7, 2017

1 Random sequence

We fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, i.e. Ω is a non-empty set, \mathcal{F} is a σ -algebra on Ω and \mathbb{P} is a probability measure on (Ω, \mathcal{F}) . We consider a sequence of real random variables $\{X_n, n \in \mathbb{N}\}$, i.e. measurable mappings $X_n : (\Omega, \mathcal{F}) \rightarrow (\mathbb{R}, \mathcal{B})$, where $\mathcal{B} = \mathcal{B}(\mathbb{R})$ is the Borel σ -algebra on \mathbb{R} .

Definition 1.1. A sequence X_1, X_2, \dots of random variables is called a *random sequence*.

Let $\mathbb{R}^{\mathbb{N}}$ be the space of all sequences of real numbers. The random sequence X_1, X_2, \dots creates a mapping $X : \Omega \rightarrow \mathbb{R}^{\mathbb{N}}$ defined by

$$X(\omega) = (X_1(\omega), X_2(\omega), \dots), \quad \omega \in \Omega. \quad (1)$$

The question is whether this mapping is measurable in some sense.

Definition 1.2. If (S_n, \mathcal{S}_n) are measurable spaces, we define the *product σ -algebra* $\bigotimes_{n=1}^{\infty} \mathcal{S}_n$ of subsets of the product space $\bigtimes_{n=1}^{\infty} S_n$ as

$$\bigotimes_{n=1}^{\infty} \mathcal{S}_n = \sigma\{A_1 \times A_2 \times \dots \times A_n \times S_{n+1} \times S_{n+2} \times \dots : A_k \in \mathcal{S}_k, n \in \mathbb{N}\}.$$

The set of the form $A_1 \times A_2 \times \dots \times A_n \times S_{n+1} \times S_{n+2} \times \dots$ is called a *finite-dimensional cylinder set*.

When $S_n = S$ and $\mathcal{S}_n = \mathcal{S}$ for all $n \in \mathbb{N}$, we use the notation $S^{\mathbb{N}} = \bigtimes_{n=1}^{\infty} S_n$ and $\mathcal{S}^{\mathbb{N}} = \bigotimes_{n=1}^{\infty} \mathcal{S}_n$.

In our situation we have $S_n = \mathbb{R}$ and $\mathcal{S}_n = \mathcal{B}$.

Proposition 1.1. *If X_1, X_2, \dots is a random sequence, then X defined in (1) is measurable with respect to $\mathcal{B}^{\mathbb{N}}$. We write $X : (\Omega, \mathcal{F}) \rightarrow (\mathbb{R}^{\mathbb{N}}, \mathcal{B}^{\mathbb{N}})$.*

Proof. Applying Definition 1.2 for $S_n = \mathbb{R}$ and $\mathcal{S}_n = \mathcal{B}$, it suffices to verify that $F = [X \in A_1 \times A_2 \times \dots \times A_n \times \mathbb{R} \times \mathbb{R} \times \dots] \in \mathcal{F}$ for any $A_1, \dots, A_n \in \mathcal{B}$ and $n \in \mathbb{N}$. However, this is obvious because $F = \bigcap_{k=1}^n [X_k \in A_k] \in \mathcal{F}$. \square

The space $\mathbb{R}^{\mathbb{N}}$ can be naturally turned into a metric space.

Definition 1.3. The distance of two real sequences $x = (x_1, x_2, \dots)$ and $y = (y_1, y_2, \dots)$ is defined as

$$d(x, y) = \sum_{n=1}^{\infty} \frac{|x_n - y_n| \wedge 1}{2^n},$$

where $a \wedge b = \min\{a, b\}$.

Remark: Note that the series in definition of d is always convergent and $d(x, y) \leq 1$ for any $x, y \in \mathbb{R}^{\mathbb{N}}$.

Proposition 1.2. (a) *The function d is a metric on $\mathbb{R}^{\mathbb{N}}$.*

(b) *The sequence $\{x^n = (x_1^n, x_2^n, \dots) \in \mathbb{R}^{\mathbb{N}}, n \in \mathbb{N}\}$ converges as $n \rightarrow \infty$ to the sequence $x = (x_1, x_2, \dots) \in \mathbb{R}^{\mathbb{N}}$ in metric d if and only if $x_j^n \xrightarrow{n \rightarrow \infty} x_j$ for all $j \in \mathbb{N}$.*

(c) *The metric space $(\mathbb{R}^{\mathbb{N}}, d)$ is separable and complete.*

(d) *For $-\infty < a_n \leq b_n < \infty$, the set $\prod_{n=1}^{\infty} [a_n, b_n]$ is a compact subset of $\mathbb{R}^{\mathbb{N}}$.*

Proof. (a), (b), (c) Exercise class.

(d) Let $\{x^n = (x_1^n, x_2^n, \dots), n \in \mathbb{N}\}$ be a sequence of elements from $[a_1, b_1] \times [a_2, b_2] \times \dots$. Since x_1^1, x_1^2, \dots is a sequence of real numbers from the compact interval $[a_1, b_1]$, there is a subsequence $x_1^{n(1,1)}, x_1^{n(1,2)}, \dots, x_1^{n(1,k)}, \dots$, that converges to some $x_1 \in [a_1, b_1]$ as $k \rightarrow \infty$. Then we can find a subsequence $\{n(2, k), k \in \mathbb{N}\}$ of $\{n(1, k), k \in \mathbb{N}\}$ such that $x_2^{n(2,k)} \xrightarrow{k \rightarrow \infty} x_2 \in [a_2, b_2]$. By induction we construct

$$\begin{aligned} x_1^{n(1,1)}, x_1^{n(1,2)}, \dots, x_1^{n(1,k)}, \dots &\xrightarrow{k \rightarrow \infty} x_1 \in [a_1, b_1], \\ x_2^{n(2,1)}, x_2^{n(2,2)}, \dots, x_2^{n(2,k)}, \dots &\xrightarrow{k \rightarrow \infty} x_2 \in [a_2, b_2], \\ &\vdots \\ x_\ell^{n(\ell,1)}, x_\ell^{n(\ell,2)}, \dots, x_\ell^{n(\ell,k)}, \dots &\xrightarrow{k \rightarrow \infty} x_\ell \in [a_\ell, b_\ell], \\ &\vdots \end{aligned}$$

so that $\{n(\ell + 1, k), k \in \mathbb{N}\}$ is a subsequence of $\{n(\ell, k), k \in \mathbb{N}\}$ for $\ell \in \mathbb{N}$. We use the diagonal selection principle and consider the sequence $\{n(k, k), k \in \mathbb{N}\}$. For any $\ell \in \mathbb{N}$ we have $\{n(k, k), k \geq \ell\} \subseteq \{n(\ell, k), k \in \mathbb{N}\}$ and hence

$$x_\ell^{n(k,k)} \xrightarrow{k \rightarrow \infty} x_\ell.$$

Using part b) we get that $\{x^{n(k,k)} = (x_1^{n(k,k)}, x_2^{n(k,k)}, \dots), k \in \mathbb{N}\}$ converges in metric d as $k \rightarrow \infty$ to the sequence $x = (x_1, x_2, \dots) \in \prod_{n=1}^{\infty} [a_n, b_n]$. This proves that $\prod_{n=1}^{\infty} [a_n, b_n]$ is compact. \square

Now let us examine what the Borel σ -algebra $\mathcal{B}(\mathbb{R}^{\mathbb{N}})$ looks like.

Theorem 1.3. *The relation $\mathcal{B}^{\mathbb{N}} = \mathcal{B}(\mathbb{R}^{\mathbb{N}})$ holds, i.e.*

$$\sigma\{A_1 \times \dots \times A_n \times \mathbb{R} \times \dots : A_1, \dots, A_n \in \mathcal{B}, n \in \mathbb{N}\} = \sigma\{U : U \subseteq \mathbb{R}^{\mathbb{N}} \text{ open set}\}.$$

Proof. The inclusion $\mathcal{B}^{\mathbb{N}} \subseteq \mathcal{B}(\mathbb{R}^{\mathbb{N}})$ is obvious. We have to show that every open set $U \subseteq \mathbb{R}^{\mathbb{N}}$ lies in the σ -algebra $\mathcal{B}^{\mathbb{N}}$. For each $x \in U$ there exists $\delta_x > 0$ such that $B(x, \delta_x) \subseteq U$, where $B(x, \delta_x) = \{y : d(y, x) < \delta_x\}$ is an open ball in $\mathbb{R}^{\mathbb{N}}$ with the centre x and radius δ_x . Therefore, $U = \bigcup_{x \in U} B(x, \delta_x)$. The metric space $\mathbb{R}^{\mathbb{N}}$ is separable (Proposition 1.2). Therefore, from an open covering of U we can select a countable subcollection (Lindelöf's covering theorem) which also covers U . We get $U = \bigcup_{k=1}^{\infty} B(x_k, \delta_{x_k})$, where $x_k \in U$. In order to finish the proof it suffices to show that $B(x, \delta) \in \mathcal{B}^{\mathbb{N}}$ for any $x \in \mathbb{R}^{\mathbb{N}}$ and $\delta > 0$. For fixed $x \in \mathbb{R}^{\mathbb{N}}$ the mapping $T : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^+$ given by $T : y \mapsto \sum_{j=1}^{\infty} 2^{-j} (|x_j - y_j| \wedge 1)$ is measurable with respect to $\mathcal{B}^{\mathbb{N}}$. Hence, $B(x, \delta) = T^{-1}((0, \delta)) \in \mathcal{B}^{\mathbb{N}}$. \square

Definition 1.4. Let E be a metric space. Then $X : (\Omega, \mathcal{F}) \rightarrow (E, \mathcal{B}(E))$ is called a *random element* with values in E .

Corollary 1.4. *A random sequence $X = (X_1, X_2, \dots)$ is a random element with values in $\mathbb{R}^{\mathbb{N}}$.*

Proof. The assertion follows from Proposition 1.1 and Theorem 1.3. \square

There exist several useful non-trivial sub- σ -algebras of the σ -algebra $\mathcal{B}(\mathbb{R}^{\mathbb{N}})$.

Definition 1.5. The mapping $p : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ is called a *finite permutation (of order n)* if there exist $n \in \mathbb{N}$ and a permutation (k_1, \dots, k_n) of the set $\{1, \dots, n\}$ such that

$$p(x_1, \dots, x_n, x_{n+1}, \dots) = (x_{k_1}, \dots, x_{k_n}, x_{n+1}, \dots), \quad (x_1, x_2, \dots) \in \mathbb{R}^{\mathbb{N}}.$$

Definition 1.6. The mapping $s : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ given by

$$s(x_1, x_2, \dots) = (x_2, x_3, \dots), \quad (x_1, x_2, \dots) \in \mathbb{R}^{\mathbb{N}},$$

is called a *shift*.

Definition 1.7. The set $T \in \mathcal{B}(\mathbb{R}^{\mathbb{N}})$ is called *terminal* if the following implication holds:

$$x = (x_1, x_2, \dots) \in T, y = (y_1, y_2, \dots) \in \mathbb{R}^{\mathbb{N}} : y_k = x_k \text{ for all but finitely many } k \in \mathbb{N} \Rightarrow y \in T.$$

We say that $T \in \mathcal{B}(\mathbb{R}^{\mathbb{N}})$ is *n-terminal* if

$$x = (x_1, x_2, \dots) \in T, y = (y_1, y_2, \dots) \in \mathbb{R}^{\mathbb{N}} : y_k = x_k \text{ for } k > n \Rightarrow y \in T.$$

Definition 1.8. Denote the following collections of sets:

- *n-symmetric sets:* $\mathcal{S}_n = \{S \in \mathcal{B}(\mathbb{R}^{\mathbb{N}}) : p(S) = S \text{ for any finite permutation } p \text{ of order } n\}$,
- *symmetric sets:* $\mathcal{S} = \{S \in \mathcal{B}(\mathbb{R}^{\mathbb{N}}) : p(S) = S \text{ for any finite permutation } p\}$,
- *shift-invariant sets:* $\mathcal{I} = \{I \in \mathcal{B}(\mathbb{R}^{\mathbb{N}}) : s^{-1}I = I\}$,
- *n-terminal sets:* $\mathcal{T}_n = \{T \in \mathcal{B}(\mathbb{R}^{\mathbb{N}}) : T \text{ n-terminal}\}$.
- *terminal sets:* $\mathcal{T} = \{T \in \mathcal{B}(\mathbb{R}^{\mathbb{N}}) : T \text{ terminal}\}$.

Proposition 1.5. (a) Any finite permutation $p : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ is a homeomorphism.

(b) Shift s is continuous mapping.

(c) The set $T \in \mathcal{B}(\mathbb{R}^{\mathbb{N}})$ is n -terminal if and only if there exists $T_n \in \mathcal{B}(\mathbb{R}^{\mathbb{N}})$ such that $T = \mathbb{R}^n \times T_n$.

(d) The collections \mathcal{I} , \mathcal{T} and \mathcal{S} are σ -algebras such that $\mathcal{I} \subset \mathcal{T}_n \subset \mathcal{S}_n$ for any $n \in \mathbb{N}$. Consequently, $\mathcal{I} \subset \mathcal{T} \subset \mathcal{S}$. All inclusions are strict, i.e. $\mathcal{I} \neq \mathcal{T}_n \neq \mathcal{S}_n$ and $\mathcal{I} \neq \mathcal{T} \neq \mathcal{S}$.

Proof. Exercise class. □

By Corollary 1.4 the random sequence $X = (X_1, X_2, \dots)$ is a random element with values in $\mathbb{R}^{\mathbb{N}}$. Hence, it has a probability distribution.

Definition 1.9. Let $X : (\Omega, \mathcal{F}) \rightarrow (E, \mathcal{B}(E))$ be a random element with values in a metric space E . Let $\mathcal{P}(E)$ denote the family of Borel probability measures on E . Define $P_X(B) = \mathbb{P}(X \in B)$ for $B \in \mathcal{B}(E)$. Then $P_X \in \mathcal{P}(E)$ is called the *probability distribution* of X .

It means that the probability distribution of a random sequence $X = (X_1, X_2, \dots)$ is a probability measure P_X on $\mathcal{B}(\mathbb{R}^{\mathbb{N}})$ defined as $P_X(B) = \mathbb{P}((X_1, X_2, \dots) \in B)$ for $B \in \mathcal{B}(\mathbb{R}^{\mathbb{N}})$. We are going to show that the distribution of X is determined by the set of finite-dimensional distributions.

Definition 1.10. We say that the set $B \in \mathcal{B}(\mathbb{R}^{\mathbb{N}})$ is *finite-dimensional* if there exist $n \in \mathbb{N}$ and $B_n \in \mathcal{B}(\mathbb{R}^n)$ such that $B = B_n \times \mathbb{R}^{\mathbb{N}}$.

Proposition 1.6. Let \mathcal{A} be the family of finite-dimensional sets. This system is an algebra that generates $\mathcal{B}(\mathbb{R}^{\mathbb{N}})$, i.e. $\sigma(\mathcal{A}) = \mathcal{B}(\mathbb{R}^{\mathbb{N}})$.

Proof. Exercise class. □

Theorem 1.7. The probability distribution of a random sequence $X = (X_1, X_2, \dots)$ is uniquely determined by the probability distribution of all random vectors (X_1, X_2, \dots, X_n) , $n \in \mathbb{N}$.

Proof. Let $X = (X_1, X_2, \dots)$ and $Y = (Y_1, Y_2, \dots)$ be random sequences satisfying $P_{(X_1, \dots, X_n)} = P_{(Y_1, \dots, Y_n)}$ for all $n \in \mathbb{N}$. We have to show that $P_X = P_Y$. Let $B = B_n \times \mathbb{R}^{\mathbb{N}}$ be a finite-dimensional set. Then

$$P_X(B) = \mathbb{P}(X \in B) = \mathbb{P}((X_1, \dots, X_n) \in B_n) = \mathbb{P}((Y_1, \dots, Y_n) \in B) = \mathbb{P}(Y \in B) = P_Y(B).$$

The measures P_X and P_Y coincide on the algebra \mathcal{A} of finite-dimensional sets. From the measure theory we know that if two finite measures coincide on some π -system (system closed under finite intersections) then they coincide on the σ -algebra generated by this π -system. Applying Proposition 1.6 we get that $P_X = P_Y$ on $\mathcal{B}(\mathbb{R}^{\mathbb{N}})$. \square

Fundamental problem: We prescribe finite-dimensional probability distributions $P_n \in \mathcal{P}(\mathbb{R}^n)$ for $n \in \mathbb{N}$. When does the random sequence $X = (X_1, X_2, \dots)$ exist such that $P_{(X_1, \dots, X_n)} = P_n$ for every $n \in \mathbb{N}$?

We easily find a necessary condition.

Definition 1.11. We say that a sequence $\{P_n \in \mathcal{P}(\mathbb{R}^n), n \in \mathbb{N}\}$ of probability distributions is *projective* if $P_{n+1}(B_n \times \mathbb{R}) = P_n(B_n)$, $B_n \in \mathcal{B}^n$, $n \in \mathbb{N}$, i.e. P_n is a marginal distribution of P_{n+1} for arbitrary $n \in \mathbb{N}$.

The distribution $P_n \in \mathcal{P}(\mathbb{R}^n)$ is uniquely determined by its distribution function

$$F_n(x_1, \dots, x_n) = P_n((-\infty, x_1] \times \dots \times (-\infty, x_n]), \quad (x_1, \dots, x_n) \in \mathbb{R}^n.$$

Therefore, we easily get the following result.

Proposition 1.8. *The system $\{P_n \in \mathcal{P}(\mathbb{R}^n), n \in \mathbb{N}\}$ is projective if and only if*

$$\lim_{x_{n+1} \rightarrow \infty} F_{n+1}(x_1, \dots, x_n, x_{n+1}) = F_n(x_1, \dots, x_n) \quad \text{for } (x_1, \dots, x_n) \in \mathbb{R}^n \text{ and } n \in \mathbb{N},$$

where F_n denotes the distribution function of P_n .

It is obvious that $\{P_n, n \in \mathbb{N}\}$ in our fundamental problem must be projective. A deep result is that this condition is also sufficient. The following two theorems are special cases of the Daniell-Kolmogorov extension theorem.

Theorem 1.9. (Daniell's extension theorem) *Let $\{P_n \in \mathcal{P}(\mathbb{R}^n), n \in \mathbb{N}\}$ be a projective family. Then there exists a random sequence $X = (X_1, X_2, \dots)$ such that $P_{(X_1, \dots, X_n)} = P_n$ for all $n \in \mathbb{N}$.*

Theorem 1.10. *Let $\{P_n \in \mathcal{P}(\mathbb{R}^n), n \in \mathbb{N}\}$ be a projective family. Then there exists a unique Borel probability measure P on $\mathbb{R}^{\mathbb{N}}$ such that*

$$P(B_n \times \mathbb{R}^{\mathbb{N}}) = P_n(B_n), \quad B_n \in \mathcal{B}^n, \quad n \in \mathbb{N}. \quad (2)$$

Theorem 1.9 is a consequence of Theorem 1.10.

Proof. (of Theorem 1.9) By Theorem 1.10 there exists $P \in \mathcal{P}(\mathbb{R}^{\mathbb{N}})$ satisfying (2). We use a canonical construction. Take $(\Omega, \mathcal{F}, \mathbb{P}) = (\mathbb{R}^{\mathbb{N}}, \mathcal{B}(\mathbb{R}^{\mathbb{N}}), P)$ and $X = \text{id}$. The projections $X_n : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}$, $n \in \mathbb{N}$, given by

$$X_n(x_1, \dots, x_n, x_{n+1}, \dots) = x_n \quad \text{for } (x_1, x_2, \dots) \in \mathbb{R}^{\mathbb{N}},$$

are continuous and hence measurable in the sense $(\mathbb{R}^{\mathbb{N}}, \mathcal{B}(\mathbb{R}^{\mathbb{N}})) \rightarrow (\mathbb{R}, \mathcal{B})$. The distribution of the random vector (X_1, \dots, X_n) is

$$P_{(X_1, \dots, X_n)}(B_n) = \mathbb{P}((X_1, \dots, X_n) \in B_n) = \mathbb{P}(X \in B_n \times \mathbb{R}^{\mathbb{N}}) = P(B_n \times \mathbb{R}^{\mathbb{N}}) = P_n(B_n), \quad B_n \in \mathcal{B}^n.$$

The random sequence $X = (X_1, X_2, \dots)$ satisfies the required property. \square

Now we prove Theorem 1.10.

Proof. (of Theorem 1.10) Let $\{P_n \in \mathcal{P}(\mathbb{R}^n), n \in \mathbb{N}\}$ be a projective system. Relation (2) defines a function P on the algebra $\mathcal{A} \subset \mathcal{B}(\mathbb{R}^{\mathbb{N}})$ of finite-dimensional sets.

We first need to verify that the definition of $P : \mathcal{A} \rightarrow [0, 1]$ is correct. Let a finite-dimensional set be expressed in two ways as $B_n \times \mathbb{R}^{\mathbb{N}} = B_m \times \mathbb{R}^{\mathbb{N}}$, where $B_n \in \mathcal{B}^n$, $B_m \in \mathcal{B}^m$ and $m > n$. Then $B_m = B_n \times \mathbb{R}^{m-n}$ and the projectivity property implies $P_m(B_m) = P_m(B_n \times \mathbb{R}^{m-n}) = P_n(B_n)$.

Next we verify that P is finitely additive on \mathcal{A} . If A and B are finite-dimensional sets, then there exist $n \in \mathbb{N}$ and $A_n, B_n \in \mathcal{B}^n$ such that $A = A_n \times \mathbb{R}^{\mathbb{N}}$ and $B = B_n \times \mathbb{R}^{\mathbb{N}}$. For disjoint sets A and B it is obvious that A_n and B_n are disjoint and $A \cup B = (A_n \cup B_n) \times \mathbb{R}^{\mathbb{N}}$. Therefore,

$$P(A \cup B) = P_n(A_n \cup B_n) = P_n(A_n) + P_n(B_n) = P(A) + P(B).$$

It remains to show that P is σ -additive probability measure on the algebra \mathcal{A} . Then by the Hahn-Kolmogorov theorem (sometimes also known as Hopf extension theorem), P can be extended to a unique probability measure \bar{P} on the σ -algebra $\sigma(\mathcal{A})$. Proposition 1.6 claims that the algebra \mathcal{A} generates $\mathcal{B}(\mathbb{R}^{\mathbb{N}})$. Thus, the extension \bar{P} is defined on $\mathcal{B}(\mathbb{R}^{\mathbb{N}})$ and has the desired properties.

Let $A^n = B_{k_n}^n \times \mathbb{R}^{\mathbb{N}}$, $n \in \mathbb{N}$, be sets from \mathcal{A} such that $A^1 \supseteq A^2 \supseteq \dots$ and $\bigcap_{n=1}^{\infty} A^n = \emptyset$ (we write shortly $A^n \searrow \emptyset$). Since A^n is a monotone sequence, we can assume that the sets $B_{k_n}^n \in \mathcal{B}^{k_n}$ are chosen so that $k_1 < k_2 < \dots$. Choose arbitrary $\varepsilon > 0$. The measure $P_{k_n} \in \mathcal{P}(\mathbb{R}^{k_n})$ is tight for any $n \in \mathbb{N}$. It means that there exists a compact set $K^n \subseteq B_{k_n}^n$ satisfying $P_{k_n}(B_{k_n}^n \setminus K^n) < \varepsilon/2^n$. We construct finite-dimensional sets $C^n = K^n \times \mathbb{R}^{\mathbb{N}} \in \mathcal{A}$. From the construction it follows that $C^n \subseteq A^n$ and hence $\bigcap_{n=1}^{\infty} C^n = \emptyset$.

We can find $m \in \mathbb{N}$ such that $\bigcap_{n=1}^m C^n = \emptyset$. For contradiction assume that $\bigcap_{n=1}^m C^n \neq \emptyset$ for all $m \in \mathbb{N}$. Then there exist sequences $x^m = (x_1^m, x_2^m, \dots) \in \mathbb{R}^{\mathbb{N}}$, $m \in \mathbb{N}$, such that $(x_1^m, \dots, x_{k_n}^m) \in K^n$ for all $n = 1, \dots, m$. So we obtained a sequence $\{x^m, m \in \mathbb{N}\}$ in $\mathbb{R}^{\mathbb{N}}$ such that every sequence $(x_\ell^1, x_\ell^2, \dots)$ is bounded in \mathbb{R} . By Proposition 1.2d, the sequence $\{x^m, m \in \mathbb{N}\}$ has a limit point $x \in \mathbb{R}^{\mathbb{N}}$. From the construction we see that $x \in \bigcap_{n=1}^{\infty} C^n = \emptyset$, which leads to the desired contradiction.

Let $m \in \mathbb{N}$ be such that $\bigcap_{n=1}^m C^n = \emptyset$. Then

$$P(A^m) = P(A^m \setminus \bigcap_{n=1}^m C^n) \leq \sum_{n=1}^m P(A^n \setminus C^n) = \sum_{n=1}^m P_{k_n}(B_{k_n}^n \setminus K^n) < \sum_{n=1}^m \frac{\varepsilon}{2^n} < \varepsilon.$$

We used the relation $A^m \setminus \bigcap_{n=1}^m C^n \subseteq \bigcup_{n=1}^m (A^n \setminus C^n)$. Hence, we have $P(A^n) \leq P(A^m) < \varepsilon$ for $n \geq m$, leading to $P(A^n) \searrow 0$ for $n \rightarrow \infty$.

If $\tilde{A}^1, \tilde{A}^2, \dots \in \mathcal{A}$ are pairwise disjoint sets such that $\bigcup_{k=1}^{\infty} \tilde{A}^k \in \mathcal{A}$, then $A^n = \bigcup_{k=n}^{\infty} \tilde{A}^k \in \mathcal{A}$ for $n \in \mathbb{N}$ and $A^n \searrow \emptyset$. We have already shown that $P(A^n) \searrow 0$ for $n \rightarrow \infty$. From this fact and finite additivity of P we can deduce that

$$P\left(\bigcup_{k=1}^{\infty} \tilde{A}^k\right) = P\left(\bigcup_{k=1}^{n-1} \tilde{A}^k\right) + P(A^n) = \sum_{k=1}^{n-1} P(\tilde{A}^k) + P(A^n) \xrightarrow{n \rightarrow \infty} \sum_{k=1}^{\infty} P(\tilde{A}^k).$$

We proved the desired σ -additivity of P , which completes the proof. \square

Recall the notion of product measure from the measure theory.

Definition 1.12. Let $Q_1, \dots, Q_n \in \mathcal{P}(\mathbb{R})$. The *product measure* $Q = \bigotimes_{k=1}^n Q_k$ is the unique probability measure in $\mathcal{P}(\mathbb{R}^n)$ satisfying the property

$$Q(B_1 \times \dots \times B_n) = Q_1(B_1) \cdots Q_n(B_n)$$

for all $B_1, \dots, B_n \in \mathcal{B}$.

For $Q_1, Q_2, \dots \in \mathcal{P}(\mathbb{R})$ there exists a unique probability measure $Q \in \mathcal{P}(\mathbb{R}^{\mathbb{N}})$ such that

$$Q(B_1 \times B_2 \times \dots \times B_n \times \mathbb{R} \times \dots) = Q_1(B_1) \cdots Q_n(B_n) = \left(\bigotimes_{k=1}^n Q_k\right)(B_1 \times \dots \times B_n)$$

for any finite-dimensional cylinder set $B_1 \times B_2 \times \dots \times B_n \times \mathbb{R} \times \dots \in \mathcal{B}(\mathbb{R}^{\mathbb{N}})$. This measure is denoted by $Q = \bigotimes_{k=1}^{\infty} Q_k$ and it is called the *infinite product measure* of Q_1, Q_2, \dots

Proposition 1.11. *The random variables X_1, \dots, X_n are independent if and only if $P_{(X_1, \dots, X_n)} = \bigotimes_{k=1}^n P_{X_k}$.*

Proof. See Probability Theory 1. □

If $\{Q_k \in \mathcal{P}(\mathbb{R}), k \in \mathbb{N}\}$ is a sequence of probability measures, it is clear that $\{P_n = \bigotimes_{k=1}^n Q_k, n \in \mathbb{N}\}$ is a projective family. Then Theorem 1.10 has the following form.

Theorem 1.12. *For an arbitrary sequence $\{Q_k \in \mathcal{P}(\mathbb{R}), k \in \mathbb{N}\}$ there exists a unique probability measure $P \in \mathcal{P}(\mathbb{R}^{\mathbb{N}})$ satisfying*

$$P(B_n \times \mathbb{R}^{\mathbb{N}}) = \left(\bigotimes_{k=1}^n Q_k \right) (B_n), \quad B_n \in \mathcal{B}^n, n \in \mathbb{N}.$$

Theorem 1.9 then can be stated in the following form.

Theorem 1.13. *For an arbitrary sequence $\{Q_k \in \mathcal{P}(\mathbb{R}), k \in \mathbb{N}\}$ there exists a sequence $X = (X_1, X_2, \dots)$ of independent random variables such that $P_{X_k} = Q_k, k \in \mathbb{N}$. Moreover, $P_X = \bigotimes_{k=1}^{\infty} Q_k$ is the infinite product measure.*

Proof. Since the system $\{P_n = \bigotimes_{k=1}^n Q_k, n \in \mathbb{N}\}$ is projective, by Theorem 1.9 there exists a random sequence $X = (X_1, X_2, \dots)$ such that $P_{(X_1, \dots, X_n)} = P_n = \bigotimes_{k=1}^n Q_k, n \in \mathbb{N}$. Hence, $P_{X_k} = Q_k$ for any $k \in \mathbb{N}$ and random variables X_1, \dots, X_n are independent for any $n \in \mathbb{N}$ by Proposition 1.11. This in turn implies that random variables X_1, X_2, \dots are independent. The equality $P_X = \bigotimes_{k=1}^{\infty} Q_k$ follows from

$$P_{(X_1, \dots, X_n)} = \bigotimes_{k=1}^n Q_k, n \in \mathbb{N}, \text{ and Definition 1.12.} \quad \square$$

Theorem 1.13 states that if we specify one-dimensional distributions Q_k , then there always exists a sequence of independent random variables X_k that have distribution $Q_k, k \in \mathbb{N}$. When the Q_k are Bernoulli distributions with parameter $p \in [0, 1]$, this gives a mathematical model for a sequence of Bernoulli trials with the probability of success p . In the case $p = 1/2$ we can proceed directly without the need of Daniell's extension theorem.

Definition 1.13. *Binary expansion* of a number $x \in (0, 1]$ is a sequence x_1, x_2, \dots of zeros and ones that contains infinitely many ones and

$$x = \sum_{k=1}^{\infty} \frac{x_k}{2^k}.$$

Binary expansion of $x = 0$ is a sequence of zeros.

Proposition 1.14. *Let X be a random variable with uniform distribution on $[0, 1]$ and let*

$$X(\omega) = \sum_{k=1}^{\infty} \frac{X_k(\omega)}{2^k} \quad (3)$$

be its binary expansion. Then X_1, X_2, \dots is a sequence of independent random variables having Bernoulli distribution with parameter $1/2$.

Conversely, if we consider a sequence of independent random variables having Bernoulli distribution with parameter $1/2$, then the random variable X defined by (3) has a uniform distribution on the interval $[0, 1]$.

Proof. Exercise class. □

We will deal with several important types of random sequences that describe the motion of a particle at times $n = 1, 2, \dots$

Definition 1.14. We say that a random sequence $X = (X_1, X_2, \dots)$ is

- *iid*, if random variables X_j , $j \in \mathbb{N}$, are independent and identically distributed,
- *n-symmetric*, if $(X_1, \dots, X_n, X_{n+1}, \dots)$ and $(X_{k_1}, \dots, X_{k_n}, X_{n+1}, \dots)$ have the same distributions for every finite permutation (k_1, \dots, k_n) of order $n \in \mathbb{N}$,
- *symmetric*, if it is *n-symmetric* for all $n \in \mathbb{N}$,
- *stationary*, if the distributions of $(X_1, \dots, X_n, X_{n+1}, \dots)$ and $(X_{n+1}, X_{n+2}, \dots)$ coincide for all $n \in \mathbb{N}$.

Examples and relations between these types of sequences are left to exercise class. Other important types are Markov chains (course Stochastic processes 1) and martingales, which we are going to study in more detail in next sections.

2 Stopping times, filtrations, and martingales

Let $X = (X_1, X_2, \dots)$ be a random sequence. It models the random motion of a particle at times $t = 1, 2, \dots$. Then the events which the particle encounters until time n are collected in the σ -algebra

$$\sigma(X_1, \dots, X_n) = \{(X_1, \dots, X_n) \in B_n, B_n \in \mathcal{B}^n\}.$$

All events are collected in σ -algebra $\sigma(X) = \{[X \in B], B \in \mathcal{B}(\mathbb{R}^{\mathbb{N}})\}$.

Proposition 2.1. *The following relation holds: $\sigma(X) = \sigma(\cup_{n=1}^{\infty} \sigma(X_1, \dots, X_n))$.*

Proof. Exercise class. □

Definition 2.1. Let (Ω, \mathcal{F}) be a measurable space and $\mathcal{F}_1 \subseteq \mathcal{F}_2 \subseteq \dots \subseteq \mathcal{F}$ be a non-decreasing sequence of σ -algebras. We say that $\{\mathcal{F}_n, n \in \mathbb{N}\}$ is a *filtration*. We denote $\mathcal{F}_\infty = \sigma(\cup_{n=1}^{\infty} \mathcal{F}_n)$.

Let $X = (X_1, X_2, \dots)$ be a sequence of random variables defined on (Ω, \mathcal{F}) and let $\{\mathcal{F}_n\}$ be a filtration satisfying $\sigma(X_1, \dots, X_n) \subseteq \mathcal{F}_n$ for all $n \in \mathbb{N}$. We say that the sequence X is *\mathcal{F}_n -adapted*.

If $\sigma(X_1, \dots, X_n) = \mathcal{F}_n$ for all $n \in \mathbb{N}$, we say that $\{\mathcal{F}_n\}$ is the *canonical filtration* of $X = (X_1, X_2, \dots)$.

Proposition 2.2. *Let $X = (X_1, X_2, \dots)$ be a random sequence and let $S = (S_1, S_2, \dots)$ be a sequence of its partial sums, i.e. $S_n = \sum_{k=1}^n X_k$, $n \in \mathbb{N}$. Then X and S have the same canonical filtration, i.e. $\sigma(X_1, \dots, X_n) = \sigma(S_1, \dots, S_n)$ for all $n \in \mathbb{N}$. Consequently, $\sigma(X) = \sigma(S)$.*

Proof. Exercise class. □

An important event occurs when a particle first enters some barrier set B .

Definition 2.2. Let $X = (X_1, X_2, \dots)$ be a random sequence. For $B \in \mathcal{B}$ denote $T_B(\omega) = \min\{n : X_n(\omega) \in B\}$, where $\min \emptyset = \infty$. We say that T_B is the *first hitting time* of the set B by the sequence X .

Note that $T_B : \Omega \rightarrow \mathbb{N} \cup \{\infty\}$ and

$$[T_B \leq n] = \bigcup_{k=1}^n [X_k \in B] \in \sigma(X_1, \dots, X_n) \subseteq \mathcal{F}, \quad n \in \mathbb{N}.$$

It means that T_B is a random variable.

Definition 2.3. The mapping $T : \Omega \rightarrow \mathbb{N} \cup \{\infty\}$ is called a *stopping time (or Markov time)* with respect to the filtration $\{\mathcal{F}_n\}$ provided that $[T \leq n] \in \mathcal{F}_n$ for all $n \in \mathbb{N}$. Shortly we speak about an *\mathcal{F}_n -stopping time (or \mathcal{F}_n -Markov time)*.

Let $X = (X_1, X_2, \dots)$ be a random sequence. A stopping time T with respect to the canonical filtration is called a *stopping time of the sequence X* , i.e. $T : \Omega \rightarrow \mathbb{N} \cup \{\infty\}$ and $[T \leq n] \in \sigma(X_1, \dots, X_n)$ for all $n \in \mathbb{N}$.

The first hitting time T_B is a stopping time of X because $[T_B \leq n] \in \sigma(X_1, \dots, X_n)$ for every $n \in \mathbb{N}$. This stopping time is a random variable that gives no information about the behaviour of X after the time T_B .

We are looking for a suitable definition of a σ -algebra that represents our information about the random sequence X up to the stopping time T .

Definition 2.4. Let $\{\mathcal{F}_n\}$ be a filtration and T be an \mathcal{F}_n -stopping time. Define

$$\mathcal{F}_T = \{F \in \mathcal{F}_\infty : F \cap [T \leq n] \in \mathcal{F}_n \forall n \in \mathbb{N}\}.$$

Then \mathcal{F}_T is a σ -algebra that is called *stopping time σ -algebra*.

Proposition 2.3. Let $\{\mathcal{F}_n\}$ be a filtration. Then T is an \mathcal{F}_n -stopping time if and only if $[T = n] \in \mathcal{F}_n$ for all $n \in \mathbb{N}$. Furthermore,

$$\mathcal{F}_T = \{F \in \mathcal{F}_\infty : F \cap [T = n] \in \mathcal{F}_n \forall n \in \mathbb{N}\}.$$

Proof. Exercise class. □

At the time $T(\omega) < \infty$ the particle is located in $X_{T(\omega)}(\omega)$. For $\omega \in \Omega$ we denote

$$X_T(\omega) = \begin{cases} X_{T(\omega)}(\omega) & \text{if } T(\omega) < \infty, \\ 0 & \text{if } T(\omega) = \infty. \end{cases}$$

If $T < \infty$ a.s. we write shortly $T \stackrel{a.s.}{<} \infty$. In that case, X_T is almost surely a value of the random sequence stopped at time T .

Proposition 2.4. (calculus for stopping times) Let $\{\mathcal{F}_n\}$ be a filtration. If S and T are \mathcal{F}_n -stopping times and $\{X_n, n \in \mathbb{N}\}$ is an \mathcal{F}_n -adapted random sequence, then

- a) T and X_T are \mathcal{F}_T -measurable random variables,
- b) $S \wedge T$, $S \vee T$ and $S + T$ are \mathcal{F}_n -stopping times,
- c) $T \wedge n$ is \mathcal{F}_n -measurable random variable for any $n \in \mathbb{N}$,
- d) $F \in \mathcal{F}_S \Rightarrow F \cap [S \leq T] \in \mathcal{F}_T$,
- e) $S \leq T \Rightarrow \mathcal{F}_S \subseteq \mathcal{F}_T$,
- f) $[S \leq T], [S = T] \in \mathcal{F}_S \cap \mathcal{F}_T$,
- g) $\mathcal{F}_S \cap \mathcal{F}_T = \mathcal{F}_{S \wedge T}$.

Proof. a), b), c) exercise.

d) According to Proposition 2.3 we have to show that $F \in \mathcal{F}_S \Rightarrow F \cap [S \leq T] \cap [T = n] \in \mathcal{F}_n$ for all $n \in \mathbb{N}$:

$$F \cap [S \leq T] \cap [T = n] = F \cap [S \leq n] \cap [T = n] \in \mathcal{F}_n.$$

This follows from $F \cap [S \leq n] \in \mathcal{F}_n$ and $[T = n] \in \mathcal{F}_n$.

e) By part d), $S \leq T$ and $F \in \mathcal{F}_S$ implies that $F = F \cap [S \leq T] \in \mathcal{F}_T$.

f) By part d), we have $[S \leq T] = \Omega \cap [S \leq T] \in \mathcal{F}_T$. If we put $\lambda = S \wedge T$, then λ is a stopping time by part b). Since $[\lambda = T] \cap [T = n] = [\lambda = n] \cap [T = n] \in \mathcal{F}_n$, we get $[\lambda = T] = [S \geq T] \in \mathcal{F}_T$ by Proposition 2.3. Altogether we have $[S \leq T], [S \geq T] \in \mathcal{F}_T$ and hence also $[S = T] = [S \leq T] \cap [S \geq T] \in \mathcal{F}_T$. The events $[S \leq T], [S \geq T], [S = T]$ belong to \mathcal{F}_S from the symmetry.

g) By part e) we get $\mathcal{F}_{S \wedge T} \subseteq \mathcal{F}_S \cap \mathcal{F}_T$. Let $F \in \mathcal{F}_S \cap \mathcal{F}_T$. Then

$$F \cap [S \wedge T \leq n] = (F \cap [T \leq S] \cap [T \leq n]) \cup (F \cap [S \leq T] \cap [S \leq n]).$$

From part f) we have $F \cap [T \leq S] \in \mathcal{F}_T$ and $F \cap [S \leq T] \in \mathcal{F}_S$. Consequently, $F \cap [T \leq S] \cap [T \leq n] \in \mathcal{F}_n$ and $F \cap [S \leq T] \cap [S \leq n] \in \mathcal{F}_n$, which leads to $F \cap [S \wedge T \leq n] \in \mathcal{F}_n$ for any $n \in \mathbb{N}$. It means that $F \in \mathcal{F}_{S \wedge T}$. □

Proposition 2.5. a) Let T be an \mathcal{F}_n -stopping time. Let $\lambda : \Omega \rightarrow \mathbb{N} \cup \{\infty\}$ be an \mathcal{F}_T -measurable random variable such that $\lambda \geq T$. Then λ is an \mathcal{F}_n -stopping time.

b) Let $X = (X_1, X_2, \dots)$ be a random sequence and T its stopping time. For $B \in \mathcal{B}$ define $\lambda = \min\{k > T : X_k \in B\}$, it is the first hitting time of B after time T . Then λ is a stopping time of X .

Proof. Exercise class. □

Definition 2.5. Let X_1, X_2, \dots be an iid random sequence and consider its partial sums $S_n = \sum_{k=1}^n X_k$, $n \in \mathbb{N}$. The sequence $\{S_n, n \in \mathbb{N}\}$ is called a *random walk*. If the random variables X_i take only values 1 and -1 , then $\{S_n\}$ is called a *simple random walk*.

Theorem 2.6. (strong Markov property of a random walk) Let $S_n = \sum_{k=1}^n X_k$ be a random walk and let $T \stackrel{a.s.}{<} \infty$ be its stopping time. Denote $R_k = S_{T+k} - S_T$ for $k \in \mathbb{N}$. Then $(R_1, R_2, \dots) \stackrel{d}{=} (S_1, S_2, \dots)$ and the sequence (R_1, R_2, \dots) is independent of the σ -algebra \mathcal{F}_T .

Proof. Consider $n \in \mathbb{N}$, $F \in \mathcal{F}_T$ and $B \in \mathcal{B}^n$. Then

$$\begin{aligned} \mathbb{P}([(R_1, \dots, R_n) \in B] \cap F) &= \sum_{k=1}^{\infty} \mathbb{P}([(R_1, \dots, R_n) \in B, T = k] \cap F) \\ &= \sum_{k=1}^{\infty} \mathbb{P}([(S_{k+1} - S_k, \dots, S_{k+n} - S_k) \in B] \cap [T = k] \cap F) \\ &= \sum_{k=1}^{\infty} \mathbb{P}((S_{k+1} - S_k, \dots, S_{k+n} - S_k) \in B) \cdot \mathbb{P}([T = k] \cap F) \\ &= \mathbb{P}((S_1, \dots, S_n) \in B) \cdot \sum_{k=1}^{\infty} \mathbb{P}([T = k] \cap F) = \mathbb{P}((S_1, \dots, S_n) \in B) \cdot \mathbb{P}(F). \end{aligned}$$

By choosing $F = \Omega$ we get $\mathbb{P}((R_1, \dots, R_n) \in B) = \mathbb{P}((S_1, \dots, S_n) \in B)$ for all $n \in \mathbb{N}$ and $B \in \mathcal{B}^n$. Applying Theorem 1.7 this in turn means that the distributions of (R_1, R_2, \dots) and (S_1, S_2, \dots) coincide. Furthermore, we have $\mathbb{P}([(R_1, \dots, R_n) \in B] \cap F) = \mathbb{P}((R_1, \dots, R_n) \in B) \cdot \mathbb{P}(F)$. Thus, (R_1, \dots, R_n) and \mathcal{F}_T are independent for any $n \in \mathbb{N}$. This is equivalent to the independence of (R_1, R_2, \dots) and \mathcal{F}_T . □

Proposition 2.7. (stationarity with respect to a stopping time) Let (X_1, X_2, \dots) be an iid random sequence and let $T \stackrel{a.s.}{<} \infty$ be its stopping time. Then $(X_{T+1}, X_{T+2}, \dots) \stackrel{d}{=} (X_1, X_2, \dots)$ and the sequence $(X_{T+1}, X_{T+2}, \dots)$ is independent of the σ -algebra \mathcal{F}_T .

Proof. Analogously as in the proof of Theorem 2.6 consider arbitrary $n \in \mathbb{N}$, $F \in \mathcal{F}_T$ and $B \in \mathcal{B}^n$. Then

$$\begin{aligned} \mathbb{P}([(X_{T+1}, \dots, X_{T+n}) \in B] \cap F) &= \sum_{k=1}^{\infty} \mathbb{P}([(X_{T+1}, \dots, X_{T+n}) \in B] \cap F \cap [T = k]) \\ &= \sum_{k=1}^{\infty} \mathbb{P}([(X_{k+1}, \dots, X_{k+n}) \in B]) \mathbb{P}(F \cap [T = k]) \\ &= \mathbb{P}([(X_1, \dots, X_n) \in B]) \sum_{k=1}^{\infty} \mathbb{P}(F \cap [T = k]) \\ &= \mathbb{P}([(X_1, \dots, X_n) \in B]) \mathbb{P}(F). \end{aligned}$$

By choosing $F = \Omega$ we get $\mathbb{P}([(X_{T+1}, \dots, X_{T+n}) \in B]) = \mathbb{P}([(X_1, \dots, X_n) \in B])$. Hence,

$$\mathbb{P}([(X_{T+1}, \dots, X_{T+n}) \in B] \cap F) = \mathbb{P}([(X_{T+1}, \dots, X_{T+n}) \in B]) \mathbb{P}(F).$$

□

Definition 2.6. Let X_1, X_2, \dots be an iid random sequence such that $\mathbb{P}(X_1 = 1) = \mathbb{P}(X_1 = -1) = 1/2$. The corresponding simple random walk $\{S_n\}$ is called the *symmetric simple random walk*.

Proposition 2.8. Consider a symmetric simple random walk $\{S_n\}$ associated with an iid random sequence (X_1, X_2, \dots) . Let $T \stackrel{a.s.}{<} \infty$ be a stopping time of this sequence. Then

$$(X_1, \dots, X_T, -X_{T+1}, -X_{T+2}, \dots) \stackrel{d}{=} (X_1, X_2, \dots).$$

Proof. The random sequence $(X_1, \dots, X_T, 0, \dots)$ is \mathcal{F}_T -measurable because for any $B \in \mathcal{B}(\mathbb{R}^{\mathbb{N}})$ and $n \in \mathbb{N}$ we have

$$[(X_1, \dots, X_T, 0, \dots) \in B] \cap [T = n] = [(X_1, \dots, X_n, 0, \dots) \in B] \cap [T = n] \in \mathcal{F}_n,$$

which implies $[(X_1, \dots, X_T, 0, \dots) \in B] \in \mathcal{F}_T$ by Proposition 2.3. Random sequences

$$(0, \dots, 0, X_{T+1}, X_{T+2}, \dots) \quad \text{and} \quad (0, \dots, 0, -X_{T+1}, -X_{T+2}, \dots)$$

have the same distribution and they are independent of \mathcal{F}_T by Proposition 2.7. Therefore, the random sequences

$$(X_1, X_2, \dots) = (X_1, \dots, X_T, 0, \dots) + (0, \dots, 0, X_{T+1}, X_{T+2}, \dots)$$

and

$$(X_1, \dots, X_T, -X_{T+1}, -X_{T+2}, \dots) = (X_1, \dots, X_T, 0, \dots) + (0, \dots, 0, -X_{T+1}, -X_{T+2}, \dots),$$

that are given as the sums of two independent sequences, have the same distribution. \square

Proposition 2.9. (reflection principle) Let $\{S_n\}$ be a symmetric simple random walk. Let T be the first hitting time of the set $\{a\}$ (for some $a \in \mathbb{N}$) by this random walk. Denote $S_k^r = 2S_{k \wedge T} - S_k$, $k \in \mathbb{N}$. Then (S_1^r, S_2^r, \dots) has the same distribution as (S_1, S_2, \dots) .

Proof. Exercise class. \square

Proposition 2.10. (maxima of symmetric simple random walk) For symmetric simple random walk $\{S_n\}$ denote $M_n = \max_{k=1, \dots, n} S_k$, $n \in \mathbb{N}$. Let T be the first hitting time of the set $\{a\}$ (for some $a \in \mathbb{N}$) by the random walk $\{S_n\}$. Then

$$\mathbb{P}(T \leq n) = \mathbb{P}(M_n \geq a) = 2\mathbb{P}(S_n \geq a) - \mathbb{P}(S_n = a) \quad \text{and} \quad \lim_{n \rightarrow \infty} \mathbb{P}(M_n \geq a) = 1.$$

Proof. Exercise class. \square

Definition 2.7. Let $H : (\Omega, \mathcal{F}) \rightarrow (E, \mathcal{B}(E))$ be a random element with values in E . We define the σ -algebra generated by H as $\sigma(H) = \{[H \in B], B \in \mathcal{B}(E)\}$. It is the smallest sub- σ -algebra $\mathcal{A} \subseteq \mathcal{F}$ such that $H : (\Omega, \mathcal{A}) \rightarrow (E, \mathcal{B}(E))$.

Definition 2.8. Consider $H : (\Omega, \mathcal{F}) \rightarrow (E, \mathcal{B}(E))$ and $T : \Omega \rightarrow \bar{\mathbb{R}}$. We say that T is H -measurable random variable if $T : (\Omega, \sigma(H)) \rightarrow (\bar{\mathbb{R}}, \bar{\mathcal{B}})$, i.e. $\sigma(T) \subseteq \sigma(H)$.

Proposition 2.11. A random variable T is H -measurable if and only if there exists $f : (E, \mathcal{B}(E)) \rightarrow (\bar{\mathbb{R}}, \bar{\mathcal{B}})$ such that $T = f(H)$.

Proof. See Probability Theory 1. \square

If T is a stopping time, then it follows that $\sigma(T) \subseteq \mathcal{F}_T$. For an example when this inclusion is sharp, consider $T = n$ for some $n \in \mathbb{N}$ such that \mathcal{F}_n is a non-trivial σ -algebra. Then $\sigma(T) = \{\emptyset, \Omega\} \subsetneq \mathcal{F}_n = \mathcal{F}_T$.

Before we get to the definition of a martingale, let us recall the definition and basic properties of the conditional expectation. We write $X \in L_1(\mathcal{F})$ for a random variable X defined on $(\Omega, \mathcal{F}, \mathbb{P})$ and satisfying $\mathbb{E}|X| < \infty$. For a σ -algebra $\mathcal{G} \subseteq \mathcal{F}$ we denote by $Y \in L_1(\mathcal{G})$ a random variable Y on $(\Omega, \mathcal{G}, \mathbb{P}|_{\mathcal{G}})$ satisfying $\mathbb{E}|Y| < \infty$.

Definition 2.9. Let $X \in L_1(\mathcal{F})$ and $\mathcal{G} \subseteq \mathcal{F}$ be a σ -algebra. A random variable $\mathbb{E}^{\mathcal{G}}X = \mathbb{E}[X|\mathcal{G}] \in L_1(\mathcal{G})$ is called *conditional expectation of X given \mathcal{G}* if for any $G \in \mathcal{G}$ we have

$$\int_G X \, d\mathbb{P} = \int_G \mathbb{E}^{\mathcal{G}}X \, d\mathbb{P}.$$

The conditional expectation $\mathbb{E}^{\mathcal{G}}X$ is \mathbb{P} -a.s. uniquely determined. We write $X \stackrel{a.s.}{=} Y$ if $\mathbb{P}(X = Y) = 1$ and $X \stackrel{a.s.}{\leq} Y$ if $\mathbb{P}(X \leq Y) = 1$.

Proposition 2.12. (calculus for conditional expectation) For $X, Y \in L_1(\mathcal{F})$ and sub- σ -algebra $\mathcal{G} \subseteq \mathcal{F}$ the following relations hold:

- a) $\mathbb{E}^{\mathcal{G}}(aX + bY + c) \stackrel{a.s.}{=} a\mathbb{E}^{\mathcal{G}}X + b\mathbb{E}^{\mathcal{G}}Y + c$ for $a, b, c \in \mathbb{R}$,
- b) $X \stackrel{a.s.}{\leq} Y \Rightarrow \mathbb{E}^{\mathcal{G}}X \stackrel{a.s.}{\leq} \mathbb{E}^{\mathcal{G}}Y$,
- c) $h : \mathbb{R} \rightarrow \mathbb{R}$ convex and $h(X) \in L_1(\mathcal{F}) \Rightarrow h(\mathbb{E}^{\mathcal{G}}X) \stackrel{a.s.}{\leq} \mathbb{E}^{\mathcal{G}}h(X)$,
- d) Y \mathcal{G} -measurable random variable and $X \cdot Y \in L_1(\mathcal{F}) \Rightarrow \mathbb{E}^{\mathcal{G}}XY \stackrel{a.s.}{=} Y \cdot \mathbb{E}^{\mathcal{G}}X$ (in particular, $X \in L_1(\mathcal{G}) \Rightarrow \mathbb{E}^{\mathcal{G}}X \stackrel{a.s.}{=} X$),
- e) $\mathcal{D} \subseteq \mathcal{F}$ sub- σ -algebra such that \mathcal{D} and $\sigma(X) \vee \mathcal{G}$ are independent $\Rightarrow \mathbb{E}[X|\mathcal{G} \vee \mathcal{D}] \stackrel{a.s.}{=} \mathbb{E}[X|\mathcal{G}]$ (notation: $\mathcal{A} \vee \mathcal{B} = \sigma(\mathcal{A} \cup \mathcal{B})$),
- f) $\mathcal{D} \subseteq \mathcal{G}$ sub- σ -algebra $\Rightarrow \mathbb{E}^{\mathcal{D}}\mathbb{E}^{\mathcal{G}}X \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{G}}\mathbb{E}^{\mathcal{D}}X \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{D}}X$ (in particular, $\mathbb{E}(\mathbb{E}^{\mathcal{G}}X) = \mathbb{E}X$),
- g) $(X, \mathbf{1}_G) \stackrel{d}{=} (Y, \mathbf{1}_G) \forall G \in \mathcal{G} \Rightarrow \mathbb{E}^{\mathcal{G}}X \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{G}}Y$.

Proof. See Probability Theory 1. □

We may also consider conditioning given the random element H with values in a metric space E . In particular, $H = (H_1, H_2, \dots)$ can be a random sequence. If $X \in L_1(\mathcal{F})$ then $\mathbb{E}[X|H] = \mathbb{E}[X|\sigma(H)]$ denotes the conditional expectation of X given H . It is a.s. uniquely determined by the conditions that $\mathbb{E}[X|H]$ is integrable H -measurable random variable and

$$\int_{[H \in B]} X \, d\mathbb{P} = \int_{[H \in B]} \mathbb{E}[X|H] \, d\mathbb{P}$$

for all $B \in \mathcal{B}(E)$. According to Proposition 2.11 there exists a Borel measurable function $f : E \rightarrow \mathbb{R}$ such that $\mathbb{E}[X|H] = f(H)$. By $f(h) = \mathbb{E}[X|H = h]$ we denote the conditional expectation of X given $H = h$.

The following two properties will later play an important role.

Proposition 2.13. Let X and Y be random elements with values in metric spaces E_1 and E_2 , respectively. Let $g : E_1 \times E_2 \rightarrow \mathbb{R}$ be a Borel measurable function such that $g(X, Y) \in L_1(\mathcal{F})$.

- (i) If $Z \in L_1(\mathcal{F})$ and (Y, Z) and X are independent, then $\mathbb{E}[Z|X, Y] \stackrel{a.s.}{=} \mathbb{E}[Z|Y]$.
- (ii) If X and Y are independent, then $\mathbb{E}[g(X, Y)|X = x] = \mathbb{E}g(x, Y)$ for P_X -a.s. $x \in E_1$.

Proof. See Probability Theory 1. □

By $X \in L_2(\mathcal{F})$ we mean that a random variable X defined on $(\Omega, \mathcal{F}, \mathbb{P})$ satisfies $\mathbb{E}X^2 < \infty$. When studying martingale differences we will need the following result.

Proposition 2.14. For $X \in L_2(\mathcal{F})$ and σ -algebra $\mathcal{G} \subset \mathcal{F}$ we have

- (i) $\mathbb{E}^{\mathcal{G}}X \in L_2(\mathcal{G})$, $\mathbb{E}(X - \mathbb{E}^{\mathcal{G}}X)^2 = \mathbb{E}X^2 - \mathbb{E}(\mathbb{E}^{\mathcal{G}}X)^2$,
- (ii) $\mathbb{E}(X - \mathbb{E}^{\mathcal{G}}X)Y = 0$ for $Y \in L_2(\mathcal{G})$,
- (iii) $\mathbb{E}(X - \mathbb{E}^{\mathcal{G}}X)^2 = \min_{Y \in L_2(\mathcal{G})} \mathbb{E}(X - Y)^2$.

Proof. See Probability Theory 1. □

The mapping $\mathbb{E}^{\mathcal{G}} : L_2(\mathcal{F}) \rightarrow L_2(\mathcal{G})$ is a projection operator in the Hilbert space L_2 . If we denote the L_2 -norm $\|X\| = \sqrt{\mathbb{E}X^2}$, then $X - Y = (X - \mathbb{E}^{\mathcal{G}}X) + (\mathbb{E}^{\mathcal{G}}X - Y)$ is the decomposition into two perpendicular summands by part (ii) of Proposition 2.14. Moreover,

$$\|X - Y\|^2 = \|X - \mathbb{E}^{\mathcal{G}}X\|^2 + \|\mathbb{E}^{\mathcal{G}}X - Y\|^2 \geq \|X - \mathbb{E}^{\mathcal{G}}X\|^2,$$

and the equality holds for $Y = \mathbb{E}^{\mathcal{G}}X$.

We will substantially improve the rule f) from Proposition 2.12 for iterated conditioning.

Proposition 2.15. *Let $\{\mathcal{F}_n\}$ be a filtration and let S and T be its stopping times. Assume that $Z \in L_1(\mathcal{F})$. Then*

(i) *the implication $(S \leq T \Rightarrow \mathbb{E}^{\mathcal{F}_S}Z = \mathbb{E}^{\mathcal{F}_{S \wedge T}}Z)$ holds a.s.,*

(ii) $\mathbb{E}^{\mathcal{F}_S}\mathbb{E}^{\mathcal{F}_T}Z \stackrel{\text{a.s.}}{=} \mathbb{E}^{\mathcal{F}_T}\mathbb{E}^{\mathcal{F}_S}Z \stackrel{\text{a.s.}}{=} \mathbb{E}^{\mathcal{F}_{S \wedge T}}Z$.

Proof. (i) We have to prove that there exists $N \in \mathcal{F}$ with the property $\mathbb{P}(N) = 0$ such that $(S(\omega) \leq T(\omega) \Rightarrow (\mathbb{E}^{\mathcal{F}_S}Z)(\omega) = (\mathbb{E}^{\mathcal{F}_{S \wedge T}}Z)(\omega))$ for $\omega \notin N$. In another words, we want to show that $\mathbf{1}_{[S \leq T]}\mathbb{E}^{\mathcal{F}_S}Z \stackrel{\text{a.s.}}{=} \mathbf{1}_{[S \leq T]}\mathbb{E}^{\mathcal{F}_{S \wedge T}}Z$. By Proposition 2.4 we have $[S \leq T] \in \mathcal{F}_{S \wedge T} \subseteq \mathcal{F}_S$. Hence, taking into account Proposition 2.12d we have to verify $\mathbb{E}^{\mathcal{F}_S}\mathbf{1}_{[S \leq T]}Z \stackrel{\text{a.s.}}{=} \mathbf{1}_{[S \leq T]}\mathbb{E}^{\mathcal{F}_{S \wedge T}}Z$. This means that we have to show

$$\int_F \mathbf{1}_{[S \leq T]}Z \, d\mathbb{P} = \int_F \mathbf{1}_{[S \leq T]}\mathbb{E}^{\mathcal{F}_{S \wedge T}}Z \, d\mathbb{P}$$

for all $F \in \mathcal{F}_S$. The last relation can be rewritten as

$$\int_{F \cap [S \leq T]} Z \, d\mathbb{P} = \int_{F \cap [S \leq T]} \mathbb{E}^{\mathcal{F}_{S \wedge T}}Z \, d\mathbb{P}. \quad (4)$$

Now it suffices to note that $F \cap [S \leq T] \in \mathcal{F}_T \cap \mathcal{F}_S = \mathcal{F}_{S \wedge T}$ by Proposition 2.4. Therefore, (4) follows from the definition of conditional expectation.

(ii) We have to show that $\mathbb{E}^{\mathcal{F}_{S \wedge T}}Z \stackrel{\text{a.s.}}{=} \mathbb{E}^{\mathcal{F}_S}(\mathbb{E}^{\mathcal{F}_T}Z)$, i.e. $\int_F \mathbb{E}^{\mathcal{F}_{S \wedge T}}Z \, d\mathbb{P} = \int_F \mathbb{E}^{\mathcal{F}_T}Z \, d\mathbb{P}$ for all $F \in \mathcal{F}_S$. For arbitrary $F \in \mathcal{F}_S$ we get

$$\int_{F \cap [S \leq T]} \mathbb{E}^{\mathcal{F}_{S \wedge T}}Z \, d\mathbb{P} = \int_{F \cap [S \leq T]} Z \, d\mathbb{P} = \int_{F \cap [S \leq T]} \mathbb{E}^{\mathcal{F}_T}Z \, d\mathbb{P}.$$

The first equality is (4) and the second equality follows from $F \cap [S \leq T] \in \mathcal{F}_T$ (see Proposition 2.4d) and the definition of conditional expectation. Similarly as in (i) we can show that $\mathbf{1}_{[T < S]}\mathbb{E}^{\mathcal{F}_{S \wedge T}}Z \stackrel{\text{a.s.}}{=} \mathbf{1}_{[T < S]}\mathbb{E}^{\mathcal{F}_T}Z$. Consequently,

$$\int_{F \cap [T < S]} \mathbb{E}^{\mathcal{F}_{S \wedge T}}Z \, d\mathbb{P} = \int_{F \cap [T < S]} \mathbb{E}^{\mathcal{F}_T}Z \, d\mathbb{P}.$$

□

The continuity of $\mathbb{E}[X \mid \mathcal{G}]$ in both arguments plays an important role. The following proposition deals with the continuity in the first argument. The continuity in the condition will be stated later.

Proposition 2.16. *Let $X_n, X \in L_1(\mathcal{F})$ and let $\mathcal{G}_n, \mathcal{G}$ be sub- σ -algebras of \mathcal{F} .*

a) *(continuity in L_1): $\mathbb{E}|X_n - X| \xrightarrow{n \rightarrow \infty} 0 \Rightarrow \mathbb{E}|\mathbb{E}^{\mathcal{G}_n}X_n - \mathbb{E}^{\mathcal{G}}X| \xrightarrow{n \rightarrow \infty} 0$,*

b) *(continuity in L_2): $\mathbb{E}(X_n - X)^2 \xrightarrow{n \rightarrow \infty} 0 \Rightarrow \mathbb{E}(\mathbb{E}^{\mathcal{G}_n}X_n - \mathbb{E}^{\mathcal{G}}X)^2 \xrightarrow{n \rightarrow \infty} 0$,*

c) *(uniform integrability): if the random sequence $\{X_n\}$ is uniformly integrable, then also the sequence $\{\mathbb{E}[X_n \mid \mathcal{G}_n]\}$ is uniformly integrable,*

d) *(monotone convergence theorem): $0 \leq X_n \xrightarrow[n \rightarrow \infty]{\text{a.s.}} X \Rightarrow 0 \leq \mathbb{E}^{\mathcal{G}_n}X_n \xrightarrow[n \rightarrow \infty]{\text{a.s.}} \mathbb{E}^{\mathcal{G}}X$,*

e) (conditional Fatou's lemma): $X_n \geq 0$, $X = \liminf_{n \rightarrow \infty} X_n \in L_1(\mathcal{F}) \Rightarrow$

$$0 \leq \mathbb{E}^{\mathcal{G}} X \leq \liminf_{n \rightarrow \infty} \mathbb{E}^{\mathcal{G}} X_n,$$

f) (dominated convergence theorem): $|X_n| \leq Y \in L_1(\mathcal{F})$, $X_n \xrightarrow[n \rightarrow \infty]{a.s.} X \Rightarrow \mathbb{E}|\mathbb{E}^{\mathcal{G}} X_n - \mathbb{E}^{\mathcal{G}} X| \xrightarrow[n \rightarrow \infty]{} 0$ and $\mathbb{E}^{\mathcal{G}} X_n \xrightarrow[n \rightarrow \infty]{a.s.} \mathbb{E}^{\mathcal{G}} X$.

Proof. a) Jensen's inequality (Proposition 2.12c) implies $|\mathbb{E}^{\mathcal{G}} X_n - \mathbb{E}^{\mathcal{G}} X| \leq \mathbb{E}^{\mathcal{G}} |X_n - X|$. Now it suffices to take expectation on both sides of the inequality.

b) Again by Jensen's inequality we have $(\mathbb{E}^{\mathcal{G}} X_n - \mathbb{E}^{\mathcal{G}} X)^2 \leq \mathbb{E}^{\mathcal{G}} (X_n - X)^2$, and taking the expectation gives the continuity in L_2 .

c) Denote $Y_n = \mathbb{E}[X_n | \mathcal{G}_n]$. Then Jensen's inequality provides $|Y_n| \leq \mathbb{E}^{\mathcal{G}_n} |X_n|$ and for the probability of a \mathcal{G}_n -measurable event $\{|Y_n| \geq c\}$ we get

$$\mathbb{P}(|Y_n| \geq c) \leq c^{-1} \int_{\{|Y_n| \geq c\}} |Y_n| d\mathbb{P} \leq c^{-1} \int_{\{|Y_n| \geq c\}} \mathbb{E}^{\mathcal{G}_n} |X_n| d\mathbb{P} \leq c^{-1} \sup_{n \in \mathbb{N}} \mathbb{E}|X_n|.$$

Hence, $\mathbb{P}(|Y_n| \geq c) \xrightarrow[c \rightarrow \infty]{} 0$ uniformly in n . Furthermore,

$$\int_{\{|Y_n| \geq c\}} |Y_n| d\mathbb{P} \leq \int_{\{|Y_n| \geq c\}} \mathbb{E}^{\mathcal{G}_n} |X_n| d\mathbb{P} = \int_{\{|Y_n| \geq c\}} |X_n| d\mathbb{P}.$$

Since the X_n have uniformly absolutely continuous integrals, the right-hand side goes to zero for $c \rightarrow \infty$ uniformly in n .

d) See Probability Theory 1.

e) By applying part d) for monotone sequence $0 \leq \inf_{k \geq n} X_k \xrightarrow[n \rightarrow \infty]{a.s.} X$ we get

$$0 \leq \mathbb{E}^{\mathcal{G}} \inf_{k \geq n} X_k \xrightarrow[n \rightarrow \infty]{a.s.} \mathbb{E}^{\mathcal{G}} X.$$

Since $\mathbb{E}^{\mathcal{G}} \inf_{k \geq n} X_k \leq \inf_{k \geq n} \mathbb{E}^{\mathcal{G}} X_k$, we have $\mathbb{E}^{\mathcal{G}} X \leq \liminf_{n \rightarrow \infty} \mathbb{E}^{\mathcal{G}} X_n$.

f) The sequences $\{Y \pm X_n\}$ are non-negative. By e) we know that

$$\mathbb{E}^{\mathcal{G}}(Y \pm X) \leq \mathbb{E}^{\mathcal{G}} Y + \liminf_{n \rightarrow \infty} \mathbb{E}^{\mathcal{G}}(\pm X_n).$$

After subtraction of $\mathbb{E}^{\mathcal{G}} Y$ we get $\pm \mathbb{E}^{\mathcal{G}} X \leq \liminf_{n \rightarrow \infty} \pm \mathbb{E}^{\mathcal{G}} X_n$, i.e. $\mathbb{E}^{\mathcal{G}} X \leq \liminf_{n \rightarrow \infty} \mathbb{E}^{\mathcal{G}} X_n$ and $\mathbb{E}^{\mathcal{G}} X \geq \limsup_{n \rightarrow \infty} \mathbb{E}^{\mathcal{G}} X_n$. Altogether,

$$\mathbb{E}^{\mathcal{G}} X \leq \liminf_{n \rightarrow \infty} \mathbb{E}^{\mathcal{G}} X_n \leq \limsup_{n \rightarrow \infty} \mathbb{E}^{\mathcal{G}} X_n \leq \mathbb{E}^{\mathcal{G}} X$$

and all the inequality signs \leq are in fact equality signs $=$. So we have proved have $\mathbb{E}^{\mathcal{G}} X_n \xrightarrow[n \rightarrow \infty]{a.s.} \mathbb{E}^{\mathcal{G}} X$.

The uniform integrability of $\{X_n\}$ and a.s. convergence imply convergence in L_1 . So the L_1 convergence of conditional expectations follows from a). □

Now we are ready to define martingales.

Definition 2.10. Let $\{\mathcal{F}_n\}$ be a filtration and let $X = (X_1, X_2, \dots)$ be a sequence of integrable random variables. We say that X is an \mathcal{F}_n -martingale if it is \mathcal{F}_n -adapted and

$$\mathbb{E}[X_{n+1} | \mathcal{F}_n] \stackrel{a.s.}{=} X_n \quad \text{for all } n \in \mathbb{N}. \quad (5)$$

In a particular case of canonical filtration $\mathcal{F}_n = \sigma(X_1, \dots, X_n)$, X is simply called a *martingale*. It satisfies

$$\mathbb{E}[X_{n+1} | X_1, X_2, \dots, X_n] \stackrel{a.s.}{=} X_n \quad \text{for all } n \in \mathbb{N}. \quad (6)$$

If $\stackrel{a.s.}{=}$ in (5) and (6) is replaced by \geq , we say that X is an \mathcal{F}_n -submartingale and *submartingale*, respectively.

If $\stackrel{a.s.}{=}$ in (5) and (6) is replaced by \leq , we say that X is an \mathcal{F}_n -supermartingale and *supermartingale*, respectively.

Remark: Obviously, $\{X_n\}$ is an \mathcal{F}_n -submartingale if and only if $\{-X_n\}$ is an \mathcal{F}_n -supermartingale.

From the definition it is clear that every martingale has constant expectation. The sequence $\{\mathbb{E}X_n\}$ is non-decreasing for a submartingale while it is non-increasing for a supermartingale.

Note that

$$(5) \iff \mathbb{E}[X_n | \mathcal{F}_k] \stackrel{a.s.}{=} X_k \quad \text{for } k \leq n,$$

$$(6) \iff \mathbb{E}[X_n | X_1, \dots, X_k] \stackrel{a.s.}{=} X_k \quad \text{for } k \leq n.$$

It is enough to use Proposition 2.12f):

$$\mathbb{E}[X_n | \mathcal{F}_k] \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{F}_k} \mathbb{E}^{\mathcal{F}_{k+1}} \dots \mathbb{E}^{\mathcal{F}_{n-1}} X_n \stackrel{a.s.}{=} X_k.$$

Similar equivalences hold for submartingales and supermartingales.

Proposition 2.17. (*stability of the martingale property*)

- (i) If a random sequence X_1, X_2, \dots is an \mathcal{F}_n -martingale, then it is also an \mathcal{G}_n -martingale for any filtration $\{\mathcal{G}_n\}$ satisfying $\sigma(X_1, \dots, X_n) \subseteq \mathcal{G}_n \subseteq \mathcal{F}_n$ for all $n \in \mathbb{N}$. In particular, each \mathcal{F}_n -martingale is a martingale.
- (ii) Let X_1, X_2, \dots be an \mathcal{F}_n -martingale and let \mathcal{D} be a σ -algebra that is independent with \mathcal{F}_∞ . Then X_1, X_2, \dots is an $(\mathcal{F}_n \vee \mathcal{D})$ -martingale.

Proof. (i) By Proposition 2.12f) we have $\mathbb{E}[X_{n+1} | \mathcal{G}_n] \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{G}_n} \mathbb{E}^{\mathcal{F}_n} X_{n+1} \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{G}_n} X_n \stackrel{a.s.}{=} X_n$.

(ii) By Proposition 2.12e) we have $\mathbb{E}[X_{n+1} | \mathcal{F}_n \vee \mathcal{D}] \stackrel{a.s.}{=} \mathbb{E}[X_{n+1} | \mathcal{F}_n] \stackrel{a.s.}{=} X_n$. □

Remark: Similar results hold for submartingales and supermartingales.

The fundamental examples of martingales are provided by sums or products of independent random variables.

Proposition 2.18. Let X_1, X_2, \dots be a sequence of independent integrable random variables. Denote $S_n = \sum_{j=1}^n X_j$ for $n \in \mathbb{N}$.

- a) If $\mathbb{E}X_n = 0$, then $\{S_n\}$ is a martingale. If $\mathbb{E}X_n \geq 0$, then $\{S_n\}$ is a submartingale. If $\mathbb{E}X_n \leq 0$, then $\{S_n\}$ is a supermartingale.
- b) If $X_n \in L_2$, $\mathbb{E}X_n = 0$ and $\mathbb{E}X_n^2 = \sigma^2$, then $M_n = S_n^2 - n\sigma^2$ is a martingale.
- c) If $\mathbb{E}X_n = 1$, then $Z_n = \prod_{j=1}^n X_j$ is a martingale.
- d) If $\mathbb{P}(X_n = -1) = q$ and $\mathbb{P}(X_n = 1) = p$, where $q = 1 - p$ and $p \in (0, 1)$, then $Y_n = (q/p)^{S_n}$ is a martingale.

Proof. a) Obviously, $S_n \in L_1$. Therefore, it suffices to realize that

$$\mathbb{E}[S_{n+1} | S_1, \dots, S_n] \stackrel{a.s.}{=} \mathbb{E}[S_{n+1} | X_1, \dots, X_n] \stackrel{a.s.}{=} \mathbb{E}[S_n + X_{n+1} | X_1, \dots, X_n] \stackrel{a.s.}{=} S_n + \mathbb{E}X_{n+1}.$$

b) The integrability of M_n follows from the assumption $X_n \in L_2$. Furthermore,

$$\mathbb{E}[S_{n+1}^2 | S_1, \dots, S_n] \stackrel{a.s.}{=} \mathbb{E}[(S_n + X_{n+1})^2 | X_1, \dots, X_n] \stackrel{a.s.}{=} S_n^2 + 2S_n\mathbb{E}X_{n+1} + \mathbb{E}X_{n+1}^2 = S_n^2 + \sigma^2,$$

which yields

$$\mathbb{E}[M_{n+1} | S_1, \dots, S_n] \stackrel{a.s.}{=} \mathbb{E}[S_{n+1}^2 | S_1, \dots, S_n] - (n+1)\sigma^2 \stackrel{a.s.}{=} S_n^2 - n\sigma^2 = M_n,$$

i.e. $\{M_n\}$ is a $\sigma(S_1, \dots, S_n)$ -martingale. Since

$$\sigma(M_1, \dots, M_n) = \sigma(S_1^2, \dots, S_n^2) \subseteq \sigma(S_1, \dots, S_n) = \sigma(X_1, \dots, X_n),$$

we conclude that $\{M_n\}$ is also a martingale by Proposition 2.17.

c), d) Exercise class. □

A convex transformation of a martingale is a submartingale.

Proposition 2.19. (i) Let X_1, X_2, \dots be an \mathcal{F}_n -martingale and let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a convex function such that $g(X_n) \in L_1$ for any $n \in \mathbb{N}$. Then $g(X_1), g(X_2), \dots$ is an \mathcal{F}_n -submartingale.

(ii) If X_1, X_2, \dots is an \mathcal{F}_n -submartingale and $g : \mathbb{R} \rightarrow \mathbb{R}$ is a convex and non-decreasing function such that $g(X_n) \in L_1$ for any $n \in \mathbb{N}$. Then $g(X_1), g(X_2), \dots$ is an \mathcal{F}_n -submartingale.

Proof. By our assumptions, $\{g(X_n)\}$ is an \mathcal{F}_n -adapted sequence of integrable random variables. From Jensen's inequality we have

$$\mathbb{E}[g(X_{n+1}) | \mathcal{F}_n] \stackrel{a.s.}{\geq} g(\mathbb{E}[X_{n+1} | \mathcal{F}_n]).$$

The right-hand side is a.s. equal to $g(X_n)$ in case (i) due to the martingale property (5) and it is a.s. greater or equal to $g(X_n)$ in case (ii) due to the submartingale property and monotonicity of g . □

Remark: In particular, $\{X_n^+\}$ is a submartingale if $\{X_n\}$ is a submartingale and $\{|Y_n|^p\}$ for $p \geq 1$ is a submartingale if $\{Y_n\}$ is a martingale.

From Proposition 2.18a we know that a random walk $\{S_n\}$ with centred steps is a martingale. Therefore, $\{S_n^2\}$ is a submartingale and by Proposition 2.18b it can be decomposed into a martingale and an increasing sequence: $S_n^2 = M_n + n\sigma^2$. It is possible to make a similar decomposition for any submartingale.

Definition 2.11. Let $\{\mathcal{F}_n\}$ be a filtration. The random sequence I_1, I_2, \dots is \mathcal{F}_n -predictable if I_n is \mathcal{F}_{n-1} -measurable for all $n \in \mathbb{N}$, where we put $\mathcal{F}_0 = \{\emptyset, \Omega\}$, i.e. I_1 is constant.

Remark: Every \mathcal{F}_n -predictable \mathcal{F}_n -martingale $\{M_n\}$ is constant a.s. because it must satisfy $M_n \stackrel{a.s.}{=} \mathbb{E}[M_{n+1} | \mathcal{F}_n] \stackrel{a.s.}{=} M_{n+1}$.

Theorem 2.20. (Doob decomposition theorem) Let $\{S_n\}$ be an \mathcal{F}_n -submartingale. Then there exist an \mathcal{F}_n -martingale $\{M_n\}$ and a non-decreasing \mathcal{F}_n -predictable sequence $\{I_n\}$ so that $S_n = M_n + I_n$, $n \in \mathbb{N}$. The summands M_n and I_n are a.s. uniquely determined under the additional condition $I_1 = 0$.

Proof. Let $\{D_n\}$ be a sequence of differences of $\{S_n\}$, i.e. $D_1 = S_1$ and $D_{n+1} = S_{n+1} - S_n$ for $n \in \mathbb{N}$. The submartingale property immediately implies $\mathbb{E}^{\mathcal{F}_n} D_{n+1} \stackrel{a.s.}{\geq} 0$, $n \in \mathbb{N}$. Put $Z_1 = 0$ and $Z_{n+1} = (\mathbb{E}^{\mathcal{F}_n} D_{n+1})^+$ for $n \in \mathbb{N}$. Then $Z_{n+1} \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{F}_n} D_{n+1}$ and $\{Z_n\}$ is \mathcal{F}_n -predictable sequence. Furthermore, we define $Y_n = D_n - Z_n$, $n \in \mathbb{N}$. Now we proceed to the cumulative sum and introduce

$$M_n = \sum_{k=1}^n Y_k, \quad I_n = \sum_{k=1}^n Z_k, \quad S_n = \sum_{k=1}^n D_k = \sum_{k=1}^n Y_k + \sum_{k=1}^n Z_k = M_n + I_n, \quad n \in \mathbb{N}.$$

We know that $I_1 = Z_1 = 0$ and $I_{n+1} = \sum_{k=1}^{n+1} Z_k$ is \mathcal{F}_n -measurable for $n \in \mathbb{N}$. It means that $\{I_n\}$ is \mathcal{F}_n -predictable sequence. Moreover, it is non-decreasing because $Z_n \geq 0$. The random sequence $\{M_n\}$ is \mathcal{F}_n -adapted as it is a difference of \mathcal{F}_n -adapted sequence $\{S_n\}$ and \mathcal{F}_n -predictable sequence $\{I_n\}$. Clearly, both D_n and Z_n are integrable. Consequently, also Y_n and M_n are integrable. We verify that $\{M_n\}$ satisfies the martingale property:

$$\mathbb{E}^{\mathcal{F}_n}(M_{n+1} - M_n) = \mathbb{E}^{\mathcal{F}_n} Y_{n+1} \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{F}_n}(D_{n+1} - \mathbb{E}^{\mathcal{F}_n} D_{n+1}) \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{F}_n} D_{n+1} - \mathbb{E}^{\mathcal{F}_n} D_{n+1} = 0,$$

i.e. $\mathbb{E}^{\mathcal{F}_n} M_{n+1} \stackrel{a.s.}{=} M_n$ and thus $\{M_n\}$ is an \mathcal{F}_n -martingale. So we found the decomposition $S_n = M_n + I_n$ into an \mathcal{F}_n -martingale and a non-decreasing \mathcal{F}_n -predictable sequence.

In order to show uniqueness assume that we have two decompositions $S_n = M_n + I_n = N_n + J_n$. Then $\bar{M}_n = M_n - N_n = J_n - I_n$ is both \mathcal{F}_n -martingale and \mathcal{F}_n -predictable sequence. According to Remark preceding this Theorem the sequence $\{\bar{M}_n\}$ is constant a.s. This constant must be zero due to the condition $I_1 = J_1 = 0$. It means that $M_n \stackrel{a.s.}{=} N_n$ and $I_n \stackrel{a.s.}{=} J_n$. \square

Definition 2.12. The sequence $\{I_n\}$ from Doob decomposition theorem is called a *compensator* of a submartingale $\{S_n\}$.

Proposition 2.21. (Martingale differences of L_2 -martingale are orthogonal in L_2) Let $\{M_n\}$ be an \mathcal{F}_n -martingale such that $M_n \in L_2$ for all $n \in \mathbb{N}$. Denote $D_1 = M_1 - \mathbb{E}M_1$ and $D_{n+1} = M_{n+1} - M_n$ for $n \in \mathbb{N}$. Then $\mathbb{E}D_n D_m = 0$ for $m \neq n$, and so $\text{var } M_n = \sum_{j=1}^n \text{var } D_j$.

Proof. First we observe that the martingale property implies $\mathbb{E}D_n = 0$ for any $n \in \mathbb{N}$. Furthermore, for $m > n$,

$$\mathbb{E}^{\mathcal{F}_n} D_m \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{F}_n} \mathbb{E}^{\mathcal{F}_{m-1}} D_m = \mathbb{E}^{\mathcal{F}_n} \mathbb{E}^{\mathcal{F}_{m-1}} (M_m - M_{m-1}) \stackrel{a.s.}{=} 0,$$

which gives

$$\mathbb{E}D_m D_n = \mathbb{E}(\mathbb{E}^{\mathcal{F}_n} D_m D_n) = \mathbb{E}(D_n \mathbb{E}^{\mathcal{F}_n} D_m) = 0.$$

It means that $\{D_n\}$ is the sequence of uncorrelated random variables and the formula for $\text{var } M_n$ follows easily from $M_n = \mathbb{E}M_1 + \sum_{j=1}^n D_j$. \square

3 Stopping theorems and maximal inequalities

Stopping problem: Let X_1, X_2, \dots be a martingale and $T_1 \leq T_2 \leq \dots$ be a sequence of its stopping times. Consider a sequence X_{T_1}, X_{T_2}, \dots given by the values of the martingale stopped at these stopping times. Is it again a martingale?

First we consider the special case $T_n = T \wedge n$, $n \in \mathbb{N}$. Then the answer is positive.

Theorem 3.1. (optional stopping theorem) Let $X = (X_1, X_2, \dots)$ be an \mathcal{F}_n -martingale (or \mathcal{F}_n -submartingale) and let T be an \mathcal{F}_n -stopping time. By stopping X at time T we obtain a random sequence $X^T = (X_{T \wedge 1}, X_{T \wedge 2}, \dots)$. This stopped sequence is an \mathcal{F}_n -martingale (or \mathcal{F}_n -submartingale).

Proof. The random variables $X_{T \wedge n}$ are $\mathcal{F}_{T \wedge n}$ -measurable (Proposition 2.4a), and so also \mathcal{F}_n -measurable ($\mathcal{F}_{T \wedge n} \subseteq \mathcal{F}_n$ by Proposition 2.4e). Therefore, X^T is \mathcal{F}_n -adapted sequence. The integrability of the random variables $X_{T \wedge n}$ follows from the simple bound

$$|X_{T \wedge n}| \leq \max_{j=1, \dots, n} |X_j| \leq \sum_{j=1}^n |X_j| \in L_1.$$

It remains to verify the martingale (or submartingale) property. We can write

$$X_{T \wedge (n+1)} = X_{T \wedge n} + D_{n+1} \mathbf{1}_{[T > n]}, \quad n \in \mathbb{N},$$

where $D_{n+1} = X_{n+1} - X_n$. Hence,

$$\mathbb{E}^{\mathcal{F}_n} X_{T \wedge (n+1)} \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{F}_n} X_{T \wedge n} + \mathbb{E}^{\mathcal{F}_n} D_{n+1} \mathbf{1}_{[T > n]} \stackrel{a.s.}{=} X_{T \wedge n} + \mathbf{1}_{[T > n]} \mathbb{E}^{\mathcal{F}_n} (X_{n+1} - X_n)$$

and the second term is zero for a martingale or greater or equal to zero for a submartingale. \square

For more general sequences of stopping times $T_1 \leq T_2 \leq \dots$ we just consider when the sequence $\{X_{T_n}\}$ is an \mathcal{F}_{T_n} -martingale. The simplest situation is for stopping times bounded by some integer $K \in \mathbb{N}$. We formulate the corresponding result for two stopping times.

Theorem 3.2. *Let X_1, X_2, \dots be an \mathcal{F}_n -martingale and let S, T be \mathcal{F}_n -stopping times such that $S \leq T \leq K < \infty$ for some $K \in \mathbb{N}$. Then $X_S, X_T \in L_1$ and*

$$\mathbb{E}^{\mathcal{F}_S} X_T \stackrel{a.s.}{=} X_S.$$

In particular, $\mathbb{E}X_T = \mathbb{E}X_S$.

If X_1, X_2, \dots is an \mathcal{F}_n -submartingale, then $X_S, X_T \in L_1$ and

$$\mathbb{E}^{\mathcal{F}_S} X_T \stackrel{a.s.}{\geq} X_S.$$

In particular, $\mathbb{E}X_T \geq \mathbb{E}X_S$.

Proof. The integrability of X_S and X_T follows similarly as in the proof of Theorem 3.1:

$$|X_T| \leq \max_{j=1, \dots, K} |X_j| \leq \sum_{j=1}^K |X_j| \in L_1.$$

First assume that $T - S \leq 1$. Then

$$\int_F (X_T - X_S) d\mathbb{P} = \sum_{j=1}^{K-1} \int_{F \cap [S=j] \cap [T>j]} (X_{j+1} - X_j) d\mathbb{P}.$$

Since $H_j = F \cap [S=j] \cap [T>j] \in \mathcal{F}_j$ for $F \in \mathcal{F}_S$, we get $\int_{H_j} (X_{j+1} - X_j) d\mathbb{P} = 0$ in case of \mathcal{F}_n -martingale $\{X_n\}$ and $\int_{H_j} (X_{j+1} - X_j) d\mathbb{P} \geq 0$ in case of \mathcal{F}_n -submartingale $\{X_n\}$. Therefore, $\int_F (X_T - X_S) d\mathbb{P} = 0$ for martingale and $\int_F (X_T - X_S) d\mathbb{P} \geq 0$ for submartingale.

In the general case we connect the times S and T by a finite chain of stopping times $V_j = (S+j) \wedge T$ satisfying $V_{j+1} - V_j \leq 1$ and $S = V_0 \leq V_1 \leq \dots \leq V_K = T$. Now we can iteratively use the result proved above for times that differ by at most 1. Then for a submartingale it follows that

$$\mathbb{E}^{\mathcal{F}_S} X_T \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{F}_{V_0}} \mathbb{E}^{\mathcal{F}_{V_1}} \dots \mathbb{E}^{\mathcal{F}_{V_{K-1}}} X_T \stackrel{a.s.}{\geq} \mathbb{E}^{\mathcal{F}_{V_0}} \mathbb{E}^{\mathcal{F}_{V_1}} \dots \mathbb{E}^{\mathcal{F}_{V_{K-2}}} X_{V_{K-1}} \stackrel{a.s.}{\geq} \dots \stackrel{a.s.}{\geq} \mathbb{E}^{\mathcal{F}_{V_0}} X_{V_1} \stackrel{a.s.}{\geq} X_S.$$

For a martingale all inequalities $\stackrel{a.s.}{\geq}$ are equalities $\stackrel{a.s.}{=}$. □

Example: Let $\{S_n\}$ be a symmetric simple random walk. Let T be its first hitting time of $\{a\}$, where $a \in \mathbb{N}$. We know that $\{S_n\}$ is a martingale and T is its stopping time that is a.s. finite. In this case we have $\mathbb{E}S_T = a > 0 = \mathbb{E}S_1$. This example shows that we cannot expect that the result of Theorem 3.2 would be valid for general unbounded stopping times.

Theorem 3.3. (i) *Let X_1, X_2, \dots be an \mathcal{F}_n -martingale and let $S \leq T \stackrel{a.s.}{<} \infty$ be \mathcal{F}_n -stopping times such that*

$$X_T \in L_1 \quad \text{and} \quad \int_{[T>n]} |X_n| d\mathbb{P} \xrightarrow[n \rightarrow \infty]{} 0. \tag{7}$$

Then $\mathbb{E}^{\mathcal{F}_S} X_T \stackrel{a.s.}{=} X_S$ (and consequently $\mathbb{E}X_T = \mathbb{E}X_S$).

(ii) *Let X_1, X_2, \dots be an \mathcal{F}_n -submartingale and let $S \leq T \stackrel{a.s.}{<} \infty$ be \mathcal{F}_n -stopping times such that*

$$X_T^+ \in L_1 \quad \text{and} \quad \int_{[T>n]} X_n^+ d\mathbb{P} \xrightarrow[n \rightarrow \infty]{} 0. \tag{8}$$

Then $\mathbb{E}^{\mathcal{F}_S} X_T \stackrel{a.s.}{\geq} X_S$ (and consequently $\mathbb{E}X_T \geq \mathbb{E}X_S$).

Proof. We are going to prove only (ii), the proof for martingales proceeds analogously.

First we realize that (8) implies $X_T \in L_1$. To see this we show that $X_T^- \in L_1$. From Theorem 3.1 we know that $\{X_{T \wedge n}\}$ is an \mathcal{F}_n -submartingale, so it must have non-decreasing expectation. In particular, $\mathbb{E}X_{T \wedge n} \geq \mathbb{E}X_1$. Thus, we obtain the following bound for the negative part

$$\begin{aligned} \mathbb{E}X_{T \wedge n}^- &= \mathbb{E}X_{T \wedge n}^+ - \mathbb{E}X_{T \wedge n} \leq \mathbb{E}X_{T \wedge n}^+ - \mathbb{E}X_1 \\ &= \mathbb{E}X_n^+ \mathbf{1}_{[T > n]} + \mathbb{E}X_T^+ \mathbf{1}_{[T \leq n]} - \mathbb{E}X_1 \leq \mathbb{E}X_n^+ \mathbf{1}_{[T > n]} + \mathbb{E}X_T^+ - \mathbb{E}X_1. \end{aligned}$$

By Fatou's lemma and (8) we have

$$\mathbb{E}X_T^- \leq \liminf_{n \rightarrow \infty} \mathbb{E}X_{T \wedge n}^- \leq \liminf_{n \rightarrow \infty} \mathbb{E}X_n^+ \mathbf{1}_{[T > n]} + \mathbb{E}X_T^+ - \mathbb{E}X_1 = \mathbb{E}X_T^+ - \mathbb{E}X_1 < \infty.$$

For a while assume that $X_S \in L_1$. We have to show that

$$\int_F X_T \, d\mathbb{P} \geq \int_F X_S \, d\mathbb{P} \quad \text{for } F \in \mathcal{F}_S. \quad (9)$$

Since $F \cap [S \leq n] = F \cap [S \leq S \wedge n] \in \mathcal{F}_{S \wedge n}$ according to Proposition 2.4d, we get from Theorem 3.2 the following relation

$$\int_{F \cap [S \leq n]} X_{T \wedge n} \, d\mathbb{P} \geq \int_{F \cap [S \leq n]} X_{S \wedge n} \, d\mathbb{P} = \int_{F \cap [S \leq n]} X_S \, d\mathbb{P}.$$

The right-hand side goes to $\int_F X_S \, d\mathbb{P}$ as $n \rightarrow \infty$ because $S < \infty$ *a.s.* and $X_S \in L_1$. The left-hand side can be rewritten as

$$\int_{F \cap [S \leq n]} X_{T \wedge n} \, d\mathbb{P} = \int_{F \cap [S \leq n] \cap [T \leq n]} X_T \, d\mathbb{P} + \int_{F \cap [S \leq n] \cap [T > n]} X_n \, d\mathbb{P}.$$

Since $S \leq T < \infty$ *a.s.* and $X_T \in L_1$, the first summand satisfies

$$\int_{F \cap [S \leq n] \cap [T \leq n]} X_T \, d\mathbb{P} = \int_{F \cap [T \leq n]} X_T \, d\mathbb{P} \xrightarrow{n \rightarrow \infty} \int_F X_T \, d\mathbb{P}.$$

For the second summand we have by (8),

$$\liminf_{n \rightarrow \infty} \int_{F \cap [S \leq n] \cap [T > n]} X_n \, d\mathbb{P} \leq \liminf_{n \rightarrow \infty} \int_{F \cap [S \leq n] \cap [T > n]} X_n^+ \, d\mathbb{P} \leq \liminf_{n \rightarrow \infty} \int_{[T > n]} X_n^+ \, d\mathbb{P} = 0.$$

Altogether,

$$\liminf_{n \rightarrow \infty} \int_{F \cap [S \leq n]} X_{T \wedge n} \, d\mathbb{P} \leq \int_F X_T \, d\mathbb{P},$$

which implies

$$\int_F X_T \, d\mathbb{P} \geq \liminf_{n \rightarrow \infty} \int_{F \cap [S \leq n]} X_{T \wedge n} \, d\mathbb{P} \geq \liminf_{n \rightarrow \infty} \int_{F \cap [S \leq n]} X_{S \wedge n} \, d\mathbb{P} = \int_F X_S \, d\mathbb{P}.$$

In other words we have shown that $\mathbb{E}^{\mathcal{F}_S} X_T \geq X_S$. This relation holds for any \mathcal{F}_n -stopping time $S \leq T$ such that $X_S \in L_1$. A particular example is $S = T \wedge k$ for arbitrary $k \in \mathbb{N}$ (integrability of $X_{T \wedge k}$ is assured by Theorem 3.2). Therefore, using Proposition 2.12c we obtain

$$X_{T \wedge k}^+ \stackrel{a.s.}{\leq} (\mathbb{E}^{\mathcal{F}_{T \wedge k}} X_T)^+ \stackrel{a.s.}{\leq} \mathbb{E}^{\mathcal{F}_{T \wedge k}} X_T^+. \quad (10)$$

Now we can get rid of assumption $X_S \in L_1$. By noticing that $[S = k] = [S = k] \cap [T \geq k] \in \mathcal{F}_k \cap \mathcal{F}_T = \mathcal{F}_{T \wedge k}$ and applying (10) we can express

$$\begin{aligned} \mathbb{E}X_S^+ &= \sum_{k=1}^{\infty} \mathbb{E}X_k^+ \mathbf{1}_{[S=k]} = \sum_{k=1}^{\infty} \mathbb{E}X_k^+ \mathbf{1}_{[S=k \leq T]} = \sum_{k=1}^{\infty} \mathbb{E}X_{T \wedge k}^+ \mathbf{1}_{[S=k \leq T]} \\ &\leq \sum_{k=1}^{\infty} \mathbb{E}X_T^+ \mathbf{1}_{[S=k \leq T]} = \sum_{k=1}^{\infty} \mathbb{E}X_T^+ \mathbf{1}_{[S=k]} = \mathbb{E}X_T^+ < \infty. \end{aligned}$$

By the same arguments as in the beginning of the proof we obtain $\mathbb{E}X_S^- < \infty$. Together it gives $X_S \in L_1$ and we have already shown that (9) holds. This completes the proof. \square

Proposition 3.4. *The condition (7) is equivalent to the condition that the stopped sequence $X^T = \{X_{T \wedge n}\}$ is uniformly integrable. Similarly, the condition (8) is equivalent to the uniform integrability of $\{X_{T \wedge n}^+\}$.*

Proof. Again we only prove the second part. If the sequence $\{X_{T \wedge n}^+\}$ is uniformly integrable, then also the sequence $X_n^+ \mathbf{1}_{[T > n]} = X_{T \wedge n}^+ \mathbf{1}_{[T > n]}$ is uniformly integrable. Moreover, it converges to zero a.s., thus also in L^1 , which means that $\mathbb{E} X_n^+ \mathbf{1}_{[T > n]} \xrightarrow{n \rightarrow \infty} 0$, i.e. $\int_{[T > n]} X_n^+ d\mathbb{P} \xrightarrow{n \rightarrow \infty} 0$. Since the limit of every uniformly integrable sequence is integrable we have

$$X_T^+ = \lim_{n \rightarrow \infty} X_{T \wedge n}^+ \mathbf{1}_{[T < \infty]} \in L_1.$$

Conversely, assume that the condition (8) holds. Then we can write $\{X_{T \wedge n}^+\}$ as the sum of two uniformly integrable sequences:

$$0 \leq X_{T \wedge n}^+ = X_n^+ \mathbf{1}_{[T > n]} + X_T^+ \mathbf{1}_{[T \leq n]}.$$

The first sequence $\{X_n^+ \mathbf{1}_{[T > n]}\}$ is uniformly integrable because it converges to zero in L_1 and the second sequence $\{X_T^+ \mathbf{1}_{[T \leq n]}\}$ is dominated by $X_T^+ \in L_1$. \square

It is natural to ask how we can check condition (7) or equivalently the uniform integrability of the sequence stopped at time T . We give several sufficient conditions that are usually much easier to verify.

Theorem 3.5. *Let X_1, X_2, \dots be an \mathcal{F}_n -martingale and let $S \leq T \stackrel{a.s.}{<} \infty$ be \mathcal{F}_n -stopping times. Consider the following conditions:*

$$\exists 0 < c < \infty : T \geq n \Rightarrow |X_n| \leq c \quad a.s., \quad (11)$$

i.e. until time T the trajectory X_1, X_2, \dots lies in the interval $[-c, c]$ a.s.;

$$(\exists 0 < c < \infty : T > n \Rightarrow |X_{n+1} - X_n| \leq c \quad a.s.) \quad \text{and} \quad \mathbb{E}T < \infty, \quad (12)$$

i.e. before time T the increments $|X_{n+1} - X_n|$ are uniformly bounded a.s. and T is integrable;

$$(\exists 0 < c < \infty : T > n \Rightarrow \mathbb{E}^{\mathcal{F}_n} |X_{n+1} - X_n| \leq c \quad a.s.) \quad \text{and} \quad \mathbb{E}T < \infty, \quad (13)$$

i.e. before time T the conditional increments are uniformly bounded a.s. and T is integrable. Then any of the conditions (11), (12) and (13) implies that

$$X_T \in L_1 \quad \text{and} \quad \mathbb{E}^{\mathcal{F}_S} X_T \stackrel{a.s.}{=} X_S.$$

Proof. The condition (12) implies the condition (13) because

$$\mathbf{1}_{[T > n]} \mathbb{E}^{\mathcal{F}_n} |X_{n+1} - X_n| \stackrel{a.s.}{=} \mathbb{E}^{\mathcal{F}_n} \mathbf{1}_{[T > n]} |X_{n+1} - X_n| \stackrel{a.s.}{\leq} c \mathbf{1}_{[T > n]}.$$

We are going to verify that both (11) and (13) are sufficient for (7). Then Theorem 3.3 finishes the proof.

First assume that (11) is satisfied. Then

$$\int_{[T > n]} |X_n| d\mathbb{P} \leq c \mathbb{P}(T > n) \xrightarrow{n \rightarrow \infty} c \mathbb{P}(T = \infty) = 0.$$

The stopped sequence $\{X_{T \wedge n}\}$ is bounded (we have $|X_{T \wedge n}| \stackrel{a.s.}{\leq} c$) so its limit X_T must be integrable. This follows from Lebesgue's dominated convergence theorem:

$$\mathbb{E}|X_T| = \lim_{n \rightarrow \infty} \mathbb{E}|X_{T \wedge n}| \leq c < \infty.$$

Now assume that condition (13) is satisfied. Define $Y_n = |X_1| + \sum_{k=1}^{n-1} |X_{k+1} - X_k|$, $n \in \mathbb{N}$. Then $|X_n| \leq Y_n$ for any $n \in \mathbb{N}$ and

$$0 \leq |X_T| \leq Y_T \stackrel{a.s.}{=} |X_1| + \sum_{k=1}^{\infty} |X_{k+1} - X_k| \mathbf{1}_{[T > k]}.$$

If we realize that condition (13) implies

$$\mathbb{E}|X_{k+1} - X_k| \mathbf{1}_{[T>k]} = \mathbb{E} \mathbf{1}_{[T>k]} \mathbb{E}^{\mathcal{F}_k} |X_{k+1} - X_k| \leq c \mathbb{P}(T > k),$$

we get

$$\mathbb{E}|X_T| \leq \mathbb{E}Y_T \leq \mathbb{E}|X_1| + \sum_{k=1}^{\infty} \mathbb{E}|X_{k+1} - X_k| \mathbf{1}_{[T>k]} \leq \mathbb{E}|X_1| + c \sum_{k=1}^{\infty} \mathbb{P}(T > k) \leq \mathbb{E}|X_1| + c\mathbb{E}T < \infty,$$

that is $X_T \in L_1$. We have also shown that $Y_T \in L_1$, which helps us verify the second part of (7):

$$\int_{[T>n]} |X_n| d\mathbb{P} \leq \int_{[T>n]} Y_n d\mathbb{P} \leq \int_{[T>n]} Y_T d\mathbb{P} \xrightarrow{n \rightarrow \infty} 0.$$

□

Remark: We can formulate similar sufficient conditions that ensure (8) for the case of a submartingale $\{X_n\}$.

Remark: In the condition (11) we are not allowed to replace $T \geq n$ with $T > n$ (see exercise class).

In applications we often consider that T is the first exit time from some bounded Borel set. Then (11) is automatically fulfilled.

Now we extend Theorem 3.3 to the setting of the Stopping problem from the beginning of Section 3.

Theorem 3.6. (optional sampling theorem)

(i) Let X_1, X_2, \dots be an \mathcal{F}_n -martingale and let $T_1 \leq T_2 \leq \dots \stackrel{a.s.}{<} \infty$ be \mathcal{F}_n -stopping times. If

$$X_{T_k} \in L_1 \quad \text{and} \quad \lim_{n \rightarrow \infty} \int_{[T_k > n]} |X_n| d\mathbb{P} = 0$$

for all $k \in \mathbb{N}$, then $(X_{T_1}, X_{T_2}, \dots)$ is an \mathcal{F}_{T_n} -martingale.

(ii) Let X_1, X_2, \dots be an \mathcal{F}_n -submartingale and let $T_1 \leq T_2 \leq \dots \stackrel{a.s.}{<} \infty$ be \mathcal{F}_n -stopping times. If

$$X_{T_k}^+ \in L_1 \quad \text{and} \quad \lim_{n \rightarrow \infty} \int_{[T_k > n]} X_n^+ d\mathbb{P} = 0,$$

for all $k \in \mathbb{N}$, then $(X_{T_1}, X_{T_2}, \dots)$ is an \mathcal{F}_{T_n} -submartingale.

Proof. The sequence $\{X_{T_n}\}$ is \mathcal{F}_{T_n} -adapted by Proposition 2.4a. The integrability of X_{T_n} is either directly assumed in case (i) or it follows from the proof of Theorem 3.3 in case (ii). The martingale or submartingale property is obtained by applying Theorem 3.3 for $T_k \leq T_{k+1}$, $k \in \mathbb{N}$. □

Remark: According to Proposition 3.4 we may equivalently rewrite the conditions in Theorem 3.6 using uniform integrability of the stopped sequences $\{X_{T_k \wedge n}, n \in \mathbb{N}\}$ and $\{X_{T_k \wedge n}^+, n \in \mathbb{N}\}$, respectively.

The following theorem provides an important application.

Theorem 3.7. (Wald's equation – general version) Let $S_n = \sum_{k=1}^n X_k$ be a random walk and let $T \in L_1$ be its stopping time. Then

a) $X_1 \in L_1 \implies S_T \stackrel{a.s.}{=} \sum_{k=1}^T X_k \in L_1$ and $\mathbb{E}S_T = \mathbb{E}T \cdot \mathbb{E}X_1$,

b) if $X_1 \in L_2$, $\mathbb{E}X_1 = 0$ and $\exists c \in (0, \infty)$ such that $(T > n \implies |S_n| \leq c \text{ a.s.}) \forall n \in \mathbb{N}$, then

$$\text{var } S_T = \mathbb{E}S_T^2 = \mathbb{E}T \cdot \mathbb{E}X_1^2 = \mathbb{E}T \cdot \text{var } X_1.$$

Proof. a) Consider $Y_n = S_n - n\mathbb{E}X_1$, $n \in \mathbb{N}$. It is a martingale by Proposition 2.18a. Furthermore,

$$\mathbb{E}^{\mathcal{F}_n} |Y_{n+1} - Y_n| = \mathbb{E}^{\mathcal{F}_n} |X_{n+1} - \mathbb{E}X_1| = \mathbb{E}|X_1 - \mathbb{E}X_1| = c < \infty,$$

where $\mathcal{F}_n = \sigma(S_1, \dots, S_n)$. It means that $\{Y_n\}$ satisfies condition (13) in Theorem 3.5. Therefore,

$$\mathbb{E}Y_T = \mathbb{E}Y_1 = 0 \implies \mathbb{E}(S_T - T\mathbb{E}X_1) = 0 \implies \mathbb{E}S_T = \mathbb{E}T \cdot \mathbb{E}X_1.$$

b) Define $M_n = S_n^2 - n\mathbb{E}X_1^2$, $n \in \mathbb{N}$. It is a martingale by Proposition 2.18b. Again we verify condition (13):

$$\begin{aligned}\mathbb{E}^{\mathcal{F}_n}|M_{n+1} - M_n| &= \mathbb{E}^{\mathcal{F}_n}|2S_nX_{n+1} + X_{n+1}^2 - \mathbb{E}X_1^2| \\ &\stackrel{a.s.}{\leq} \mathbb{E}^{\mathcal{F}_n}2|S_n||X_{n+1}| + \mathbb{E}^{\mathcal{F}_n}X_{n+1}^2 + \mathbb{E}X_1^2 \stackrel{a.s.}{=} 2|S_n|\mathbb{E}|X_1| + 2\mathbb{E}X_1^2.\end{aligned}$$

Hence,

$$\mathbf{1}_{[T > n]}\mathbb{E}^{\mathcal{F}_n}|M_{n+1} - M_n| \stackrel{a.s.}{\leq} 2\mathbf{1}_{[T > n]}|S_n|\mathbb{E}|X_1| + 2\mathbb{E}X_1^2 \stackrel{a.s.}{\leq} 2c\mathbb{E}|X_1| + 2\mathbb{E}X_1^2 < \infty.$$

Theorem 3.5 gives

$$0 = \mathbb{E}M_1 = \mathbb{E}M_T = \mathbb{E}(S_T^2 - T\mathbb{E}X_1^2) = \mathbb{E}S_T^2 - \mathbb{E}T \cdot \mathbb{E}X_1^2,$$

which implies $\mathbb{E}S_T^2 = \mathbb{E}T \cdot \mathbb{E}X_1^2$. □

We notice that Theorem 3.7 has a simpler version.

Proposition 3.8. (Wald's equation – basic version) *Let $S_n = \sum_{k=1}^n X_k$ be a random walk and let $T \in L_1$ be independent of $\{S_n\}$. Then*

a) $X_1 \in L_1 \implies \mathbb{E}S_T = \mathbb{E}T \cdot \mathbb{E}X_1$,

b) $X_1 \in L_2, \mathbb{E}X_1 = 0 \implies \text{var } S_T = \mathbb{E}S_T^2 = \mathbb{E}T \cdot \mathbb{E}X_1^2 = \mathbb{E}T \cdot \text{var } X_1$.

Proof. Denote $\mathcal{G}_n = \sigma(S_1, \dots, S_n) \vee \sigma(T)$. Then T is a \mathcal{G}_n -stopping time and $Y_n = S_n - n\mathbb{E}X_1$ is a \mathcal{G}_n -martingale (by Proposition 2.17). The proof of part a) follows in the same way as the proof of part a) in Theorem 3.7. Another possibility is to use the independence and directly calculate $\mathbb{E}S_T$ (see Probability Theory 1).

Part b) could be shown directly:

$$\mathbb{E}S_T^2 = \mathbb{E} \sum_{k=1}^{\infty} \mathbf{1}_{[T=k]} S_k^2 = \sum_{k=1}^{\infty} \mathbb{P}(T=k) \mathbb{E}S_k^2 = \mathbb{E}X_1^2 \sum_{k=1}^{\infty} k \mathbb{P}(T=k) = \mathbb{E}T \cdot \mathbb{E}X_1^2.$$

□

The application of optional stopping theorem and optional sampling theorem to the simple random walk is left to exercise class.

Definition 3.1. We say that the random walk $\{S_n\}$ with steps $\{X_n\}$ is *nontrivial* if $\mathbb{P}(X_1 \neq 0) > 0$.

A nontrivial random walk has one of the following properties (see exercise class): 1. $S_n \xrightarrow[n \rightarrow \infty]{a.s.} \infty$, 2. $S_n \xrightarrow[n \rightarrow \infty]{a.s.} -\infty$, 3. $\limsup_{n \rightarrow \infty} S_n = \infty$ and $\liminf_{n \rightarrow \infty} S_n = -\infty$ a.s. In particular, if T^B is the first exit time of a nontrivial random walk from a bounded Borel set $B \in \mathcal{B}(\mathbb{R})$, then $T^B \stackrel{a.s.}{<} \infty$. Moreover, T^B has all moments finite.

Theorem 3.9. *Let $T^B = \min\{n : S_n \notin B\}$ be the first exit time of a nontrivial random walk $\{S_n\}$ from a bounded Borel set $B \in \mathcal{B}(\mathbb{R})$. Then $\mathbb{E}(T^B)^r < \infty$ for all $r \in \mathbb{N}$.*

Proof. We already mentioned that $T^B \stackrel{a.s.}{<} \infty$. Denote by d the diameter of the set B and find n_0 such that $P = \mathbb{P}(|S_{n_0}| > d) > 0$. Consider a new random walk $\{Z_k, k \in \mathbb{N}\}$ defined by $Z_k = S_{kn_0}$, $k \in \mathbb{N}$. This random walk moves faster than $\{S_n\}$. Let λ be the first exit of $\{Z_n\}$ from B . Then $\lambda \stackrel{a.s.}{<} \infty$ and $T^B \leq n_0\lambda$. Furthermore, $\mathbb{P}(\lambda = k) \leq \mathbb{P}(S_{n_0} \in B, |S_{(j+1)n_0} - S_{jn_0}| \leq d \text{ for } j = 1, \dots, k-2) \leq (1-P)^{k-2}$ for $k > 2$. For $k = 2$ we also have $\mathbb{P}(\lambda = k) \leq (1-P)^{k-2} = 1$. Therefore, $\mathbb{E}\lambda^r = \sum_{k=1}^{\infty} k^r \mathbb{P}(\lambda = k) \leq \mathbb{P}(\lambda = 1) + \sum_{k=2}^{\infty} k^r (1-P)^{k-2} < \infty$. □

Using Theorem 3.9 we get a transparent variant of Theorem 3.7.

Theorem 3.10. Let $\{S_n\}$ be a nontrivial random walk and let T be its first exit time from some bounded Borel set. Then

$$a) X_1 \in L_1 \implies S_T \stackrel{a.s.}{=} \sum_{k=1}^T X_k \in L_1 \text{ and } \mathbb{E}S_T = \mathbb{E}T \cdot \mathbb{E}X_1,$$

$$b) X_1 \in L_2, \mathbb{E}X_1 = 0 \implies S_T \in L_2 \text{ and } \text{var } S_T = \mathbb{E}S_T^2 = \mathbb{E}T \cdot \mathbb{E}X_1^2 = \mathbb{E}T \cdot \text{var } X_1.$$

Proof. It is a consequence of Theorem 3.7 as T is an integrable stopping time which satisfies $T > n \implies |S_n| \leq c$ a.s. \square

The following theorem gives another application of the stopping theory.

Theorem 3.11. (supermartingale goes bankrupt forever) Let X_1, X_2, \dots be a non-negative supermartingale. Consider $T = \min\{n : X_n = 0\}$, where $\min \emptyset = \infty$. Then the implication $(T < \infty \implies X_{T+k} = 0 \text{ for } k \in \mathbb{N})$ holds a.s.

Proof. If $T \stackrel{a.s.}{=} \infty$ there is nothing to prove. Let $\mathbb{P}(T < \infty) > 0$ and define $n_0 = \min\{n \in \mathbb{N} : \mathbb{P}(T \leq n) > 0\}$. Take $n \geq n_0$ and $k \in \mathbb{N}$. Denote $T_n = T \wedge n$. Then $T_n \leq T_n + k \leq n + k$ are stopping times and the variant of Theorem 3.2 for supermartingale yields

$$\mathbb{E}^{\mathcal{F}_{T_n}} X_{T_n+k} \stackrel{a.s.}{\leq} X_{T_n}.$$

Since $[T \leq n] \in \mathcal{F}_T \cap \mathcal{F}_n = \mathcal{F}_{T_n}$ by Proposition 2.4, we have

$$0 \leq \int_{[T \leq n]} X_{T+k} d\mathbb{P} = \int_{[T \leq n]} X_{T_n+k} d\mathbb{P} \leq \int_{[T \leq n]} X_{T_n} d\mathbb{P} = \int_{[T \leq n]} X_T d\mathbb{P} = 0.$$

It means that $X_{T+k} \mathbf{1}_{[T \leq n]} \stackrel{a.s.}{=} 0$. Taking the limit as $n \rightarrow \infty$ we obtain $X_{T+k} \mathbf{1}_{[T < \infty]} \stackrel{a.s.}{=} 0$. In other words, there exists a \mathbb{P} -null set N_k (i.e. $\mathbb{P}(N_k) = 0$) such that $X_{T+k}(\omega) \mathbf{1}_{[T(\omega) < \infty]} = 0$ for $\omega \notin N_k$. From this it follows that the sequence $(X_{T+k}(\omega) \mathbf{1}_{[T(\omega) < \infty]}, k \in \mathbb{N})$ is equal to the null sequence for $\omega \notin N = \bigcup_{k=1}^{\infty} N_k$, where $\mathbb{P}(N) = 0$. \square

The theory of martingales was developed by a distinguished American mathematician J. L. Doob (1910–2004). We formulate two maximal inequalities that are named after him. However, first we prove the following lemma.

Lemma 3.12. Let X_1, X_2, \dots be an \mathcal{F}_n -submartingale. If we denote $M_n = \max_{k=1, \dots, n} X_k$ for $n \in \mathbb{N}$, then

$$\mathbb{P}(M_n \geq a) \leq a^{-1} \int_{[M_n \geq a]} X_n^+ d\mathbb{P} \leq a^{-1} \mathbb{E}X_n^+$$

for any $a > 0$.

Proof. Define $F_k = [X_1 < a, \dots, X_{k-1} < a, X_k \geq a] \in \mathcal{F}_k$, $k \in \mathbb{N}$. Proposition 2.19 implies that X_1^+, X_2^+, \dots is an \mathcal{F}_n -submartingale. Hence, for any $n \in \mathbb{N}$ we can write

$$a\mathbb{P}(M_n \geq a) = a\mathbb{P}\left(\bigcup_{k=1}^n F_k\right) = a \sum_{k=1}^n \mathbb{P}(F_k) \leq \sum_{k=1}^n \int_{F_k} X_k^+ d\mathbb{P} \leq \sum_{k=1}^n \int_{F_k} X_n^+ d\mathbb{P} = \int_{[M_n \geq a]} X_n^+ d\mathbb{P} \leq \mathbb{E}X_n^+.$$

\square

Theorem 3.13. (Doob's maximal inequalities) Let X_1, X_2, \dots be a martingale or a non-negative submartingale. Then for all $n \in \mathbb{N}$ we have

1.

$$\mathbb{P}\left(\max_{k=1, \dots, n} |X_k| \geq a\right) \leq a^{-p} \mathbb{E}|X_n|^p, \quad \text{for } p \geq 1 \text{ and } a > 0,$$

2. (Doob's L_p inequality)

$$\mathbb{E}\left(\max_{k=1, \dots, n} |X_k|\right)^p \leq \left(\frac{p}{p-1}\right)^p \mathbb{E}|X_n|^p, \quad \text{for } p > 1.$$

Proof. Let us fix $n \in \mathbb{N}$. Obviously we can assume that $X_n \in L_p$, otherwise the right-hand side is infinite and the inequalities hold trivially. If $\{X_n\}$ is a martingale or a non-negative submartingale, then $\{|X_n|^p\}$ with $p \geq 1$ is a non-negative submartingale by Proposition 2.19. Applying Lemma 3.12 to this submartingale gives the first inequality.

Let $Y = \max_{k=1, \dots, n} |X_k|$ and fix $p > 1$. Then

$$\mathbb{E}Y^p \leq \sum_{k=1}^n \mathbb{E}|X_k|^p \leq n\mathbb{E}|X_n|^p < \infty.$$

The expectation of a non-negative random variable is obtained by integrating its complementary distribution function. Therefore,

$$\mathbb{E}Y^p = \int_0^\infty \mathbb{P}(Y^p > t) dt = \int_0^\infty \mathbb{P}(Y^p > a^p) p a^{p-1} da.$$

If we use Lemma 3.12 for a submartingale $|X_1|, |X_2|, \dots$ and Fubini's theorem, we get the bound

$$\begin{aligned} \mathbb{E}Y^p &\leq \int_0^\infty p a^{p-1} a^{-1} \int_{[Y \geq a]} |X_n| d\mathbb{P} da = \mathbb{E} \int_0^\infty p a^{p-2} |X_n| \mathbf{1}_{[Y \geq a]} da \\ &= p \mathbb{E}|X_n| \frac{Y^{p-1}}{p-1} = \frac{p}{p-1} \mathbb{E}|X_n| Y^{p-1}. \end{aligned}$$

By Hölder's inequality,

$$\mathbb{E}|X_n| Y^{p-1} \leq (\mathbb{E}|X_n|^p)^{\frac{1}{p}} (\mathbb{E}Y^p)^{\frac{p-1}{p}}.$$

Therefore,

$$\mathbb{E}Y^p \leq \frac{p}{p-1} (\mathbb{E}|X_n|^p)^{\frac{1}{p}} (\mathbb{E}Y^p)^{\frac{p-1}{p}}.$$

Doob's L_p inequality is trivially satisfied for $\mathbb{E}Y^p = 0$. If $\mathbb{E}Y^p > 0$ we can divide both sides by $(\mathbb{E}Y^p)^{\frac{p-1}{p}}$ and obtain

$$(\mathbb{E}Y^p)^{1/p} \leq \frac{p}{p-1} (\mathbb{E}|X_n|^p)^{1/p},$$

which is equivalent to the formulation of Doob's L_p inequality. \square

Doob's maximal inequality implies the classical maximal inequalities for independent random variables.

Theorem 3.14. (Kolmogorov's inequality) *Let X_1, X_2, \dots be independent random variables with zero expectation and finite variance and let $S_n = X_1 + \dots + X_n$. Then*

$$\mathbb{P}\left(\max_{k=1, \dots, n} |S_k| \geq a\right) \leq a^{-2} \mathbb{E}S_n^2 = a^{-2} \sum_{k=1}^n \mathbb{E}X_k^2, \quad a > 0.$$

Proof. Since $\{S_n\}$ is a martingale, it is enough to use the first Doob's maximal inequality for $p = 2$. \square

4 Submartingale convergence

Definition 4.1. Consider real numbers $a < b$ and a finite sequence $y^{(n)} = (y_1, \dots, y_n)$ of real numbers. We denote the *number of upcrossings of (a, b)* by

$$y^{(n)} \uparrow_a^b = \text{card}\{(s, t) : 1 \leq s < t \leq n, \{y_{s+1}, \dots, y_{t-1}\} \subseteq (a, b) \subseteq [y_s, y_t]\},$$

where we let $[c, d] = \emptyset$ for $c > d$. Analogously, the *number of downcrossings of (a, b)* is

$$y^{(n)} \downarrow_a^b = \text{card}\{(s, t) : 1 \leq s < t \leq n, \{y_{s+1}, \dots, y_{t-1}\} \subseteq (a, b) \subseteq [y_t, y_s]\}.$$

For an infinite sequence $y = (y_1, y_2, \dots)$ we put

$$y \uparrow_a^b = \lim_{n \rightarrow \infty} y^{(n)} \uparrow_a^b \quad \text{and} \quad y \downarrow_a^b = \lim_{n \rightarrow \infty} y^{(n)} \downarrow_a^b.$$

Remark: Clearly, $y \uparrow_a^b = (-y) \downarrow_{-b}^{-a}$ and $y \downarrow_a^b - 1 \leq y \uparrow_a^b \leq y \downarrow_a^b + 1$. If $X = (X_1, X_2, \dots)$ is a random sequence, then $X \uparrow_a^b$ is a random variable with values in $\mathbb{N} \cup \{0, \infty\}$.

Proposition 4.1. *Let $X = (X_1, X_2, \dots)$ be a random sequence.*

- (i) *There exists a random variable X^* (with values in $\mathbb{R} \cup \{-\infty, \infty\}$) such that $X_n \xrightarrow[n \rightarrow \infty]{\text{a.s.}} X^*$ if and only if $\mathbb{P}(X \uparrow_a^b < \infty) = 1$ for each $a, b \in \mathbb{R} : a < b$.*
- (ii) *There exists a real random variable Y such that $X_n \xrightarrow[n \rightarrow \infty]{\text{a.s.}} Y$ if and only if $\mathbb{P}(\sup_{n \in \mathbb{N}} |X_n| < \infty) = 1$ and $\mathbb{P}(X \uparrow_a^b < \infty) = 1$ for each $a, b \in \mathbb{R} : a < b$.*

Proof. Both implications from left to right are obvious.

Assume that $\mathbb{P}(X \uparrow_a^b < \infty) = 1$ for each $a, b \in \mathbb{R} : a < b$ and consider the random variables

$$X^* = \limsup_{n \rightarrow \infty} X_n \quad \text{and} \quad X_* = \liminf_{n \rightarrow \infty} X_n.$$

Then

$$\mathbb{P}(X_* < X^*) \leq \sum_{a, b \in \mathbb{Q} : a < b} \mathbb{P}(X_* < a < b < X^*) \leq \sum_{a, b \in \mathbb{Q} : a < b} \mathbb{P}(X \uparrow_a^b = \infty) = 0,$$

and so $X_n \xrightarrow[n \rightarrow \infty]{\text{a.s.}} X^*$.

If we moreover assume that $\sup_{n \in \mathbb{N}} |X_n| \stackrel{\text{a.s.}}{<} \infty$, then $|X^*| \leq \sup_{n \in \mathbb{N}} |X_n| \stackrel{\text{a.s.}}{<} \infty$ and we can take $Y = X^* \mathbf{1}_{\{|X^*| < \infty\}}$. \square

Theorem 4.2. (Doob's upcrossing inequality) *Let $\{X_n\}$ be an \mathcal{F}_n -submartingale. Then for $n \in \mathbb{N}$ and $a, b \in \mathbb{R}$ such that $a < b$,*

$$\mathbb{E}X^{(n) \uparrow_a^b} \leq \frac{\mathbb{E}(X_n - a)^+ - \mathbb{E}(X_1 - a)^+}{b - a} \leq \frac{\mathbb{E}(X_n - a)^+}{b - a},$$

where $X^{(n)} = (X_1, \dots, X_n)$.

Proof. Consider a sequence $Z_n = (X_n - a)^+$, $n \in \mathbb{N}$. By Proposition 2.19 we know that $\{Z_n\}$ is an \mathcal{F}_n -submartingale. Define \mathcal{F}_n -stopping times $\tau_0 = 1$, $\nu_j = \min\{k \geq \tau_{j-1} : Z_k = 0\} \wedge n$, $\tau_j = \min\{k \geq \nu_j : Z_k \geq b - a\} \wedge n$. From the definition we see that $\tau_0 \leq \nu_1 \leq \tau_1 \leq \nu_2 \leq \dots \leq n$. Moreover, $\nu_j < n$ implies $\nu_j < \tau_j$. Similarly, $\tau_j < n$ implies $\tau_j < \nu_{j+1}$. Therefore, there exists $m \in \mathbb{N}$ such that $\tau_m = \nu_m = n$. We can write

$$Z_n - Z_1 = \sum_{j=1}^m (Z_{\nu_j} - Z_{\tau_{j-1}}) + \sum_{j=1}^m (Z_{\tau_j} - Z_{\nu_j}) \geq \sum_{j=1}^m (Z_{\nu_j} - Z_{\tau_{j-1}}) + (b - a)Z^{(n) \uparrow_0^{b-a}},$$

where $Z^{(n)} = (Z_1, \dots, Z_n)$. According to Theorem 3.2 for bounded times $\tau_{j-1} \leq \nu_j$ we have $\mathbb{E}Z_{\nu_j} \geq \mathbb{E}Z_{\tau_{j-1}}$ for any j . Hence,

$$\mathbb{E}(Z_n - Z_1) \geq (b - a)\mathbb{E}Z^{(n) \uparrow_0^{b-a}}.$$

Finally, it suffices to use the definition of $\{Z_n\}$ which yields

$$\mathbb{E}(X_n - a)^+ - \mathbb{E}(X_1 - a)^+ \geq (b - a)\mathbb{E}X^{(n) \uparrow_a^b},$$

because $Z^{(n) \uparrow_0^{b-a}} = X^{(n) \uparrow_a^b}$. \square

Remark: The statement for an \mathcal{F}_n -supermartingale $\{X_n\}$ has the form

$$\mathbb{E}X^{(n) \downarrow_a^b} \leq \frac{\mathbb{E}(b - X_n)^+ - \mathbb{E}(b - X_1)^+}{b - a} \leq \frac{\mathbb{E}(b - X_n)^+}{b - a},$$

as follows from the relation $y^{(n) \uparrow_a^b} = (-y)^{(n) \downarrow_{-b}^{-a}}$.

Theorem 4.3. (Doob's submartingale convergence theorem) Let $\{X_n\}$ be an \mathcal{F}_n -submartingale that satisfies $\sup_{n \in \mathbb{N}} \mathbb{E}X_n^+ < \infty$. Then there exists a random variable $X_\infty \in L_1$ such that $X_n \xrightarrow[n \rightarrow \infty]{\text{a.s.}} X_\infty$ and

$$\begin{aligned} \mathbb{E}X_\infty^+ &\leq \liminf_{n \rightarrow \infty} \mathbb{E}X_n^+ \leq \sup_{n \in \mathbb{N}} \mathbb{E}X_n^+ < \infty \\ \mathbb{E}X_\infty^- &\leq \liminf_{n \rightarrow \infty} \mathbb{E}X_n^- \leq \sup_{n \in \mathbb{N}} \mathbb{E}X_n^+ - \mathbb{E}X_1 < \infty. \end{aligned}$$

Proof. For $a < b$, $X \uparrow_a^b$ is the limit of a non-decreasing non-negative sequence $X^{(n)} \uparrow_a^b$, where $X^{(n)} = (X_1, \dots, X_n)$. Hence, from Lévi's monotone convergence theorem and Theorem 4.2 we obtain

$$\begin{aligned} \mathbb{E}X \uparrow_a^b &= \lim_{n \rightarrow \infty} \mathbb{E}X^{(n)} \uparrow_a^b \leq \liminf_{n \rightarrow \infty} \frac{\mathbb{E}(X_n - a)^+}{b - a} \\ &\leq \liminf_{n \rightarrow \infty} \frac{\mathbb{E}X_n^+ + a^-}{b - a} \leq \frac{\sup_{n \in \mathbb{N}} \mathbb{E}X_n^+ + a^-}{b - a} < \infty. \end{aligned}$$

By Proposition 4.1 there exists a random variable X_∞ such that $X_n \xrightarrow[n \rightarrow \infty]{\text{a.s.}} X_\infty$. Positive and negative part are continuous functions, thus also $X_n^+ \xrightarrow[n \rightarrow \infty]{\text{a.s.}} X_\infty^+$ and $X_n^- \xrightarrow[n \rightarrow \infty]{\text{a.s.}} X_\infty^-$. From Fatou's lemma we get

$$\begin{aligned} \mathbb{E}X_\infty^+ &= \mathbb{E} \liminf_{n \rightarrow \infty} X_n^+ \leq \liminf_{n \rightarrow \infty} \mathbb{E}X_n^+ \leq \sup_{n \in \mathbb{N}} \mathbb{E}X_n^+ < \infty, \\ \mathbb{E}X_\infty^- &= \mathbb{E} \liminf_{n \rightarrow \infty} X_n^- \leq \liminf_{n \rightarrow \infty} (\mathbb{E}X_n^+ - \mathbb{E}X_n) \leq \sup_{n \in \mathbb{N}} \mathbb{E}X_n^+ - \mathbb{E}X_1 < \infty. \end{aligned}$$

We used that the submartingale $\{X_n\}$ satisfies $\mathbb{E}X_n \geq \mathbb{E}X_1$ for any $n \in \mathbb{N}$. Altogether, $\mathbb{E}|X_\infty| = \mathbb{E}X_\infty^+ + \mathbb{E}X_\infty^- < \infty$. \square

Remark: Similarly, every \mathcal{F}_n -supermartingale satisfying $\sup_{n \in \mathbb{N}} \mathbb{E}X_n^- < \infty$ has an integrable a.s.-limit. As special cases we have the following statements:

1. A submartingale bounded from above has an integrable a.s.-limit.
2. A supermartingale bounded from below (e.g. a non-negative supermartingale) has an integrable a.s.-limit.
3. Each martingale that is bounded either from above or from below has an integrable a.s.-limit.

Corollary 4.4. Let $\{X_n\}$ be a sequence of independent random variables such that $\mathbb{E}X_n = 0$ and

$$\sup_{n \in \mathbb{N}} \mathbb{E} \left| \sum_{k=1}^n X_k \right| < \infty.$$

Then $\sum_{k=1}^\infty X_k$ is a.s.-summable and $\sum_{k=1}^\infty X_k \in L_1$.

Proof. Define $S_n = \sum_{k=1}^n X_k$, $n \in \mathbb{N}$. Then $\{S_n\}$ is a martingale and we assume that $\sup_{n \in \mathbb{N}} \mathbb{E}|S_n| < \infty$. So it suffices to use Theorem 4.3. \square

Remark: Recall that the condition $\sum_{n=1}^\infty \text{var } X_n < \infty$ is sufficient for the summability of $\sum_{n=1}^\infty (X_n - \mathbb{E}X_n)$ a.s., in probability and in L_2 (see Probability Theory 1). In this case we have an improvement for a.s.-summability. The condition $\sum_{n=1}^\infty \text{var } X_n < \infty$ implies $\sup_{n \in \mathbb{N}} \mathbb{E}|S_n| < \infty$ because

$$\mathbb{E}|S_n| \leq \sqrt{\mathbb{E}S_n^2} = \sqrt{\sum_{k=1}^n \text{var } X_k} \leq \sqrt{\sum_{k=1}^\infty \text{var } X_k}.$$

Definition 4.2. Let (\dots, X_{-2}, X_{-1}) be a random sequence indexed by negative integers. Let $\dots \subseteq \mathcal{F}_{-2} \subseteq \mathcal{F}_{-1}$ be a non-decreasing sequence of σ -algebras (filtration). Assume that $X_{-n} \in L_1$ for any $n \in \mathbb{N}$ and $\sigma(\dots, X_{-n-1}, X_{-n}) \subseteq \mathcal{F}_{-n}$. We say that the sequence $\{X_{-n}\}$ is an \mathcal{F}_{-n} -martingale if

$$\mathbb{E}[X_{-n} \mid \mathcal{F}_{-(n+1)}] \stackrel{\text{a.s.}}{=} X_{-(n+1)} \quad \text{for all } n \in \mathbb{N}.$$

If $\mathcal{F}_{-n} = \sigma(\dots, X_{-n-1}, X_{-n})$, then we speak about a *backwards martingale*. Analogously we define \mathcal{F}_{-n} -submartingale and \mathcal{F}_{-n} -supermartingale. We denote $\mathcal{F}_{-\infty} = \bigcap_{n=1}^\infty \mathcal{F}_{-n}$.

An example of a backwards martingale can be constructed from the random walk.

Lemma 4.5. *Let $\{X_n\}$ be an iid random sequence of integrable random variables. Define*

$$Z_{-n} = \frac{1}{n} \sum_{k=1}^n X_k, \quad n \in \mathbb{N}.$$

Then $\{Z_{-n}\}$ is a backwards martingale.

Proof. By the triangle inequality, $\mathbb{E}|Z_{-n}| \leq \mathbb{E}|X_1| < \infty$. Denote $\mathcal{F}_{-n} = \sigma(\dots, Z_{-n-1}, Z_{-n})$ and consider the sequence $M_{-n} = \mathbb{E}[X_1 | \mathcal{F}_{-n}]$, $n \in \mathbb{N}$. From symmetry we have $\mathbb{E}[X_1 | \mathcal{F}_{-n}] \stackrel{a.s.}{=} \dots \stackrel{a.s.}{=} \mathbb{E}[X_n | \mathcal{F}_{-n}]$. Hence,

$$\mathbb{E}[Z_{-n+1} | \mathcal{F}_{-n}] \stackrel{a.s.}{=} \frac{1}{n-1} \sum_{k=1}^{n-1} \mathbb{E}[X_k | \mathcal{F}_{-n}] \stackrel{a.s.}{=} M_{-n}.$$

However, we can also write

$$Z_{-n} \stackrel{a.s.}{=} \mathbb{E}[Z_{-n} | \mathcal{F}_{-n}] \stackrel{a.s.}{=} \frac{1}{n} \sum_{k=1}^n \mathbb{E}[X_k | \mathcal{F}_{-n}] \stackrel{a.s.}{=} M_{-n}$$

for any $n \in \mathbb{N}$. □

We can formulate the analogy of Theorem 4.3 for backwards submartingales.

Theorem 4.6. (Doob's backwards submartingale convergence theorem) *Let $\{X_{-n}\}$ be an \mathcal{F}_{-n} -submartingale. Then there exists a random variable $X_{-\infty}$ (with values in $\mathbb{R} \cup \{-\infty, \infty\}$) such that $X_{-n} \xrightarrow[n \rightarrow \infty]{a.s.} X_{-\infty}$. The limiting random variable $X_{-\infty}$ is integrable provided that $\sup_{n \in \mathbb{N}} \mathbb{E}X_{-n}^- < \infty$.*

Proof. Denote $X = (\dots, X_{-2}, X_{-1})$ and $X^{(-n)} = (X_{-n}, \dots, X_{-1})$ for $n \in \mathbb{N}$. Similarly as in the proof of Theorem 4.3 we obtain

$$\mathbb{E}X \uparrow_a^b = \lim_{n \rightarrow \infty} \mathbb{E}X^{(-n)} \uparrow_a^b \leq \frac{\mathbb{E}(X_{-1} - a)^+}{b - a} < \infty.$$

We have used the version of Doob's upcrossing inequality (Theorem 4.2) for backwards submartingales. The existence of the a.s.-limit $X_{-\infty}$ now follows from Proposition 4.1. In order to show $X_{-\infty} \in L_1$ apply Fatou's lemma for both positive and negative part,

$$\mathbb{E}X_{-\infty}^+ \leq \liminf_{n \rightarrow \infty} \mathbb{E}X_{-n}^+ \leq \mathbb{E}X_{-1}^+ < \infty$$

and

$$\mathbb{E}X_{-\infty}^- \leq \liminf_{n \rightarrow \infty} \mathbb{E}X_{-n}^- \leq \sup_{n \in \mathbb{N}} \mathbb{E}X_{-n}^-.$$

□

Theorem 4.7. (convergence of uniformly integrable (sub)martingale)

- a) *Let $\{X_n\}$ be a uniformly integrable \mathcal{F}_n -submartingale (or \mathcal{F}_n -martingale), then there exists a random variable $X_\infty \in L_1$ such that both $X_n \xrightarrow[n \rightarrow \infty]{a.s.} X_\infty$ and $X_n \xrightarrow[n \rightarrow \infty]{L_1} X_\infty$. Furthermore, $\mathbb{E}[X_\infty | \mathcal{F}_n] \stackrel{a.s.}{\geq} X_n$ (or $\mathbb{E}[X_\infty | \mathcal{F}_n] \stackrel{a.s.}{=} X_n$) for all $n \in \mathbb{N}$.*
- b) *Let $\{X_{-n}\}$ be a uniformly integrable \mathcal{F}_{-n} -submartingale (or \mathcal{F}_{-n} -martingale), then there exists a random variable $X_{-\infty} \in L_1$ such that both $X_{-n} \xrightarrow[n \rightarrow \infty]{a.s.} X_{-\infty}$ and $X_{-n} \xrightarrow[n \rightarrow \infty]{L_1} X_{-\infty}$. Furthermore, $\mathbb{E}[X_{-n} | \mathcal{F}_{-\infty}] \stackrel{a.s.}{\geq} X_{-\infty}$ (or $\mathbb{E}[X_{-n} | \mathcal{F}_{-\infty}] \stackrel{a.s.}{=} X_{-\infty}$) for all $n \in \mathbb{N}$.*

Proof. a) The assumption of uniform integrability implies uniformly bounded moments and the assumption of Doob's submartingale convergence theorem (Theorem 4.3) is fulfilled: $\sup_{n \in \mathbb{N}} \mathbb{E}X_n^+ \leq \sup_{n \in \mathbb{N}} \mathbb{E}|X_n| < \infty$. Hence, there exists a random variable $X_\infty \in L_1$ such that $X_n \xrightarrow[n \rightarrow \infty]{a.s.} X_\infty$. We obtain the L_1 convergence from the property of uniformly integrable sequences. For fixed integer numbers $n \leq m$ we have $X_n \leq^{a.s.} \mathbb{E}[X_m | \mathcal{F}_n]$. By the L_1 continuity of conditional expectation (Proposition 2.16a), $\mathbb{E}[X_m | \mathcal{F}_n]$ converges to $\mathbb{E}[X_\infty | \mathcal{F}_n]$ as $m \rightarrow \infty$. The L_1 convergence implies convergence in probability which in turn implies the existence of a subsequence that converges a.s. For this subsequence $\{m_k\}$ we already know that $X_n \leq^{a.s.} \mathbb{E}[X_{m_k} | \mathcal{F}_n]$. This inequality is preserved when passing to the limit. Hence, $X_n \leq^{a.s.} \mathbb{E}[X_\infty | \mathcal{F}_n]$.

b) Theorem 4.6 and uniform integrability ensure the existence of a random variable $X_{-\infty} \in L_1$ such that $X_{-n} \xrightarrow[n \rightarrow \infty]{a.s.} X_{-\infty}$ and $X_{-n} \xrightarrow[n \rightarrow \infty]{L_1} X_{-\infty}$. Then the L_1 continuity of conditional expectation (Proposition 2.16a) yields $\mathbb{E}[X_{-m} | \mathcal{F}_{-\infty}] \xrightarrow[m \rightarrow \infty]{L_1} \mathbb{E}[X_{-\infty} | \mathcal{F}_{-\infty}] = X_{-\infty}$. From $X_{-m} \leq^{a.s.} \mathbb{E}[X_{-n} | \mathcal{F}_{-m}]$ for $m \geq n$ we get by conditioning with the σ -algebra $\mathcal{F}_{-\infty}$ the relation $\mathbb{E}[X_{-m} | \mathcal{F}_{-\infty}] \leq^{a.s.} \mathbb{E}[X_{-n} | \mathcal{F}_{-\infty}]$. The left-hand side goes in L_1 to $X_{-\infty}$ as $m \rightarrow \infty$ and the inequality is preserved when passing to the limit. \square

Corollary 4.8. *If $\{X_n\}$ is an \mathcal{F}_n -adapted random sequence, then $\{X_n\}$ is a uniformly integrable \mathcal{F}_n -martingale if and only if there exists $X_\infty \in L_1$ such that $X_n \stackrel{a.s.}{=} \mathbb{E}[X_\infty | \mathcal{F}_n]$.*

Proof. The implication from left to right follows from Theorem 4.7. Conversely, the sequence given by $X_n \stackrel{a.s.}{=} \mathbb{E}[X_\infty | \mathcal{F}_n]$ is an \mathcal{F}_n -martingale (see exercise class) that is uniformly integrable (Proposition 2.16c). \square

Convergence theorems imply that conditional expectations are continuous in the condition.

Proposition 4.9. *Let $Y \in L_1$ and $\cdots \subseteq \mathcal{F}_{-2} \subseteq \mathcal{F}_{-1} \subseteq \mathcal{F}$, $\mathcal{F}_1 \subseteq \mathcal{F}_2 \subseteq \cdots \subseteq \mathcal{F}$ be non-decreasing sequences of σ -algebras. Then*

- a) $\mathbb{E}[Y | \mathcal{F}_n] \xrightarrow[n \rightarrow \infty]{} \mathbb{E}[Y | \mathcal{F}_\infty]$ both a.s. and in L_1 ,
- b) $\mathbb{E}[Y | \mathcal{F}_{-n}] \xrightarrow[n \rightarrow \infty]{} \mathbb{E}[Y | \mathcal{F}_{-\infty}]$ both a.s. and in L_1 .

Proof. a) We know that $Y_n = \mathbb{E}[Y | \mathcal{F}_n]$ is uniformly integrable \mathcal{F}_n -martingale (Corollary 4.8). By Theorem 4.7 there exists $Y_\infty \in L_1$ so that $Y_n \xrightarrow[n \rightarrow \infty]{} Y_\infty$ both a.s. and in L_1 and $Y_n \stackrel{a.s.}{=} \mathbb{E}[Y_\infty | \mathcal{F}_n]$. We are going to show that $\mathbb{E}[Y | \mathcal{F}_\infty] \stackrel{a.s.}{=} Y_\infty$. Denote $Y^* = \limsup_{n \rightarrow \infty} \mathbb{E}[Y | \mathcal{F}_n]$. For $F \in \mathcal{F}_n$ we have

$$\int_F Y \, d\mathbb{P} = \int_F Y_n \, d\mathbb{P} = \int_F Y_\infty \, d\mathbb{P} = \int_F Y^* \, d\mathbb{P}.$$

The first equality follows from $Y_n = \mathbb{E}[Y | \mathcal{F}_n]$, the second equality from $Y_n \stackrel{a.s.}{=} \mathbb{E}[Y_\infty | \mathcal{F}_n]$ and the last equality from $Y_\infty \stackrel{a.s.}{=} Y^*$. Thus, we have verified the relation

$$\int_F Y \, d\mathbb{P} = \int_F Y^* \, d\mathbb{P}$$

for arbitrary $F \in \cup_{n=1}^\infty \mathcal{F}_n$, where $\cup_{n=1}^\infty \mathcal{F}_n$ is an algebra that generates \mathcal{F}_∞ . Moreover, Y^* is \mathcal{F}_∞ -measurable, which yields $\mathbb{E}[Y | \mathcal{F}_\infty] \stackrel{a.s.}{=} Y^*$.

b) A random sequence $Y_{-n} = \mathbb{E}[Y | \mathcal{F}_{-n}]$ is uniformly integrable backwards martingale (exercise class). By Theorem 4.7 there exists a random variable $Y_{-\infty} \in L_1$ such that $Y_{-n} \xrightarrow[n \rightarrow \infty]{} Y_{-\infty}$ both a.s. and in L_1 and $Y_{-n} \stackrel{a.s.}{=} \mathbb{E}[Y_{-\infty} | \mathcal{F}_{-n}]$. Then for arbitrary $F \in \mathcal{F}_{-\infty}$ we have

$$\int_F Y_{-\infty} \, d\mathbb{P} = \int_F Y_{-n} \, d\mathbb{P} = \int_F Y \, d\mathbb{P}.$$

In other words, $Y_{-\infty} \stackrel{\text{a.s.}}{=} \mathbb{E}[Y \mid \mathcal{F}_{-\infty}]$. □

Theorem 4.10. (submartingale converges or explodes) *Let $\{X_n\}$ be a submartingale. Denote $Y_k = X_{k+1} - X_k$ for $k \in \mathbb{N}$. If $(\sup_{n \in \mathbb{N}} Y_n)^+ \in L_1$, then there exists a random variable X_∞ such that $X_n(\omega) \xrightarrow[n \rightarrow \infty]{} X_\infty(\omega)$ for a.a. $\omega \in \Omega$ with the property $\sup_{n \in \mathbb{N}} X_n(\omega) < \infty$.*

Proof. For $k \in \mathbb{N}$ denote the stopping time $\tau_k = \min\{n \in \mathbb{N} : X_n \geq k\}$. We fix $k \in \mathbb{N}$. Optional stopping theorem (Theorem 3.1) states that $(X_{n \wedge \tau_k})$ is a submartingale. We distinguish the following three cases:

1. $\tau_k = 1 \Rightarrow X_{n \wedge \tau_k} = X_1$,
2. $1 < \tau_k \leq n \Rightarrow X_{n \wedge \tau_k} = X_{\tau_k} = X_{\tau_k - 1} + Y_{\tau_k - 1} \leq k + \sup_{n \in \mathbb{N}} Y_n$,
3. $\tau_k > n \Rightarrow X_{n \wedge \tau_k} = X_n < k$.

Combining all three cases we have

$$X_{n \wedge \tau_k}^+ \leq X_1^+ + k + \left(\sup_{n \in \mathbb{N}} Y_n \right)^+ \in L_1.$$

It means that $\sup_{n \in \mathbb{N}} \mathbb{E}X_{n \wedge \tau_k}^+ < \infty$ and we can apply Doob's submartingale convergence theorem (Theorem 4.3). Therefore, there exists a random variable $X^{(k)} \in L_1$ such that $X_{n \wedge \tau_k} \xrightarrow[n \rightarrow \infty]{\text{a.s.}} X^{(k)}$. Then for $A_k = [\tau_k = \infty] = [\sup_{n \in \mathbb{N}} X_n < k]$,

$$X_n \mathbf{1}_{A_k} = X_{n \wedge \tau_k} \mathbf{1}_{A_k} \xrightarrow[n \rightarrow \infty]{\text{a.s.}} X^{(k)} \mathbf{1}_{A_k}.$$

The events A_k form a non-decreasing sequence and their limit for $k \rightarrow \infty$ is $A = [\sup_{n \in \mathbb{N}} X_n < \infty]$. Furthermore, $X^{(k)} \mathbf{1}_{A_k} \stackrel{\text{a.s.}}{=} X^{(l)} \mathbf{1}_{A_k}$ for $l \geq k$. If we put $X_\infty = X^{(1)} \mathbf{1}_{A_1} + X^{(2)} \mathbf{1}_{A_2 \setminus A_1} + \dots$, then $X_n \mathbf{1}_A \xrightarrow[n \rightarrow \infty]{\text{a.s.}} X_\infty$. □

5 Limit theorems for martingale differences

Definition 5.1. Let $\{M_n\}$ be a martingale. Put $M_0 = \mathbb{E}M_1$ and define $D_n = M_n - M_{n-1}$ for $n \in \mathbb{N}$. Then $\{D_n\}$ is called a *martingale difference sequence* of the martingale $\{M_n\}$. If $\{M_n\}$ is an \mathcal{F}_n -martingale we speak about an \mathcal{F}_n -*martingale difference sequence*.

Remark: Equivalently we can define an \mathcal{F}_n -martingale difference sequence as the sequence satisfying $\mathbb{E}[D_n \mid \mathcal{F}_{n-1}] = 0$ for $n \in \mathbb{N}$, where we let $\mathcal{F}_0 = \{\emptyset, \Omega\}$.

Theorem 5.1. (summability of martingale differences) *Let $\{D_n\}$ be a martingale difference sequence of the martingale $\{M_n\}$ that satisfies $M_n \in L_2$ for each $n \in \mathbb{N}$. If $\sum_{n=1}^{\infty} \text{var } D_n < \infty$ the series $\sum_{n=1}^{\infty} D_n$ is summable both a.s. and in L_2 , i.e. the martingale $M_n - \mathbb{E}M_1$ converges both a.s. and in L_2 .*

Proof. According to Proposition 2.21 the random variables D_n are uncorrelated. Recall that from Probability Theory 1 we know that centred uncorrelated random variables are summable in L_2 if and only if the sum of their variances is finite. In order to get a.s.-summability we verify the assumption of Theorem 4.3:

$$\begin{aligned} \mathbb{E}|M_n - \mathbb{E}M_1| &\leq \sqrt{\mathbb{E}(M_n - \mathbb{E}M_1)^2} = \sqrt{\mathbb{E} \left(\sum_{k=1}^n D_k \right)^2} = \sqrt{\text{var} \sum_{k=1}^n D_k} \\ &= \sqrt{\sum_{k=1}^n \text{var } D_k} \leq \sqrt{\sum_{k=1}^{\infty} \text{var } D_k} < \infty, \end{aligned}$$

and so $\sup_{n \in \mathbb{N}} \mathbb{E}|M_n - \mathbb{E}M_1| < \infty$. □

Theorem 5.2. (strong law of large numbers for martingale differences) Let $\{D_n\}$ be a martingale difference sequence of the martingale $\{M_n\}$ that satisfies $M_n \in L_2$ for each $n \in \mathbb{N}$. Let $0 < b_n \nearrow_{n \rightarrow \infty} \infty$ be a real sequence. If $\sum_{n=1}^{\infty} b_n^{-2} \text{var } D_n < \infty$, then

$$\frac{1}{b_n} \sum_{k=1}^n D_k = \frac{M_n - \mathbb{E}M_1}{b_n} \xrightarrow[n \rightarrow \infty]{} 0 \quad \text{both a.s. and in } L_2.$$

Proof. The sequence $\left\{\frac{D_n}{b_n}\right\}$ is also a martingale difference sequence and it satisfies the assumption of Theorem 5.1. Therefore, the series $\sum_{n=1}^{\infty} b_n^{-1} D_n$ is a.s.-summable. The a.s.-convergence of $b_n^{-1} \sum_{k=1}^n D_k$ follows from Kronecker's lemma which says that if $\sum_{n=1}^{\infty} a_n < \infty$ and $0 < b_n \nearrow_{n \rightarrow \infty} \infty$, then $\frac{1}{b_n} \sum_{k=1}^n a_k b_k \xrightarrow[n \rightarrow \infty]{} 0$. To show convergence in L_2 we apply Kronecker's lemma as well:

$$\mathbb{E} \left(\frac{1}{b_n} \sum_{k=1}^n D_k \right)^2 = \frac{1}{b_n^2} \sum_{k=1}^n \mathbb{E} D_k^2 = \frac{1}{b_n^2} \sum_{k=1}^n \text{var } D_k \xrightarrow[n \rightarrow \infty]{} 0.$$

□

Theorem 5.3. (central limit theorem for martingale differences) Let $\{D_n\}$ be an \mathcal{F}_n -martingale difference sequence of the martingale $\{M_n\}$. Assume that for each $n \in \mathbb{N}$ we have

1. $\mathbb{E}[D_n^2 | \mathcal{F}_{n-1}] = 1$,
2. $\mathbb{E}[|D_n|^3 | \mathcal{F}_{n-1}] \leq K < \infty$,

where $\mathcal{F}_0 = \{\emptyset, \Omega\}$. Then

$$\frac{1}{\sqrt{n}} \sum_{k=1}^n D_k = \frac{1}{\sqrt{n}} (M_n - \mathbb{E}M_1) \xrightarrow[n \rightarrow \infty]{d} N(0, 1).$$

Proof. Define

$$\varphi_{n,k}(t) = \mathbb{E} \left[\exp \left\{ it \frac{D_k}{\sqrt{n}} \right\} \middle| \mathcal{F}_{k-1} \right], \quad k = 1, \dots, n, \quad n \in \mathbb{N}.$$

From Taylor's expansion we get

$$\exp \left\{ it \frac{D_k}{\sqrt{n}} \right\} = 1 + it \frac{D_k}{\sqrt{n}} - \frac{t^2 D_k^2}{2n} - it^3 \frac{\Delta_k^3}{6n^{3/2}},$$

where Δ_k is a random variable such that $0 \leq \Delta_k \leq D_k$. Applying conditional expectation on both sides we obtain

$$\varphi_{n,k}(t) = 1 + \frac{it}{\sqrt{n}} \mathbb{E}[D_k | \mathcal{F}_{k-1}] - \frac{t^2}{2n} \mathbb{E}[D_k^2 | \mathcal{F}_{k-1}] - \frac{it^3}{6n^{3/2}} \mathbb{E}[\Delta_k^3 | \mathcal{F}_{k-1}],$$

which by our assumptions can be simplified to

$$\varphi_{n,k}(t) = 1 - \frac{t^2}{2n} - \frac{it^3}{6n^{3/2}} \mathbb{E}[\Delta_k^3 | \mathcal{F}_{k-1}].$$

For $p = 1, \dots, n$ we have

$$\begin{aligned} \mathbb{E} \exp \left\{ it \frac{M_p}{\sqrt{n}} \right\} &= \mathbb{E} \left[\exp \left\{ it \frac{M_{p-1}}{\sqrt{n}} \right\} \exp \left\{ it \frac{D_p}{\sqrt{n}} \right\} \right] = \mathbb{E} \left[\exp \left\{ it \frac{M_{p-1}}{\sqrt{n}} \right\} \mathbb{E} \left[\exp \left\{ it \frac{D_p}{\sqrt{n}} \right\} \middle| \mathcal{F}_{p-1} \right] \right] \\ &= \mathbb{E} \left[\exp \left\{ it \frac{M_{p-1}}{\sqrt{n}} \right\} \varphi_{n,p}(t) \right] = \mathbb{E} \left[\exp \left\{ it \frac{M_{p-1}}{\sqrt{n}} \right\} \left(1 - \frac{t^2}{2n} - \frac{it^3}{6n^{3/2}} \mathbb{E}[\Delta_p^3 | \mathcal{F}_{p-1}] \right) \right]. \end{aligned}$$

Consequently,

$$\mathbb{E} \exp \left\{ it \frac{M_p}{\sqrt{n}} \right\} - \left(1 - \frac{t^2}{2n} \right) \mathbb{E} \exp \left\{ it \frac{M_{p-1}}{\sqrt{n}} \right\} = -\frac{it^3}{6n^{3/2}} \mathbb{E} \left[\exp \left\{ it \frac{M_{p-1}}{\sqrt{n}} \right\} \mathbb{E}[\Delta_p^3 | \mathcal{F}_{p-1}] \right].$$

Since $|\Delta_p| \leq |D_p|$ and the conditional absolute third moments are bounded, it follows that

$$\left| \mathbb{E} \exp \left\{ it \frac{M_p}{\sqrt{n}} \right\} - \left(1 - \frac{t^2}{2n} \right) \mathbb{E} \exp \left\{ it \frac{M_{p-1}}{\sqrt{n}} \right\} \right| \leq K \frac{|t|^3}{6n^{3/2}}. \quad (14)$$

Let us fix $t \in \mathbb{R}$. For a sufficiently large n (namely, $n \geq t^2/2$) we have $0 \leq 1 - \frac{t^2}{2n} \leq 1$. Therefore, the left-hand side of (14) is not going to increase by multiplication with $\left(1 - \frac{t^2}{2n}\right)^{n-p}$. It means that

$$\left| \left(1 - \frac{t^2}{2n} \right)^{n-p} \mathbb{E} \exp \left\{ it \frac{M_p}{\sqrt{n}} \right\} - \left(1 - \frac{t^2}{2n} \right)^{n-p+1} \mathbb{E} \exp \left\{ it \frac{M_{p-1}}{\sqrt{n}} \right\} \right| \leq K \frac{|t|^3}{6n^{3/2}}.$$

We apply the triangle inequality to the identity

$$\begin{aligned} & \mathbb{E} \exp \left\{ it \frac{M_n - \mathbb{E}M_1}{\sqrt{n}} \right\} - \left(1 - \frac{t^2}{2n} \right)^n \\ &= \exp \left\{ -it \frac{\mathbb{E}M_1}{\sqrt{n}} \right\} \sum_{p=1}^n \left[\left(1 - \frac{t^2}{2n} \right)^{n-p} \mathbb{E} \exp \left\{ it \frac{M_p}{\sqrt{n}} \right\} - \left(1 - \frac{t^2}{2n} \right)^{n-p+1} \mathbb{E} \exp \left\{ it \frac{M_{p-1}}{\sqrt{n}} \right\} \right] \end{aligned}$$

and obtain (for $n \geq t^2/2$)

$$\left| \mathbb{E} \exp \left\{ it \frac{M_n - \mathbb{E}M_1}{\sqrt{n}} \right\} - \left(1 - \frac{t^2}{2n} \right)^n \right| \leq nK \frac{|t|^3}{6n^{3/2}} = K \frac{|t|^3}{6\sqrt{n}}.$$

As the right-hand side tends to zero and $\left(1 - \frac{t^2}{2n}\right)^n$ tends to $e^{-t^2/2}$ for $n \rightarrow \infty$, we have

$$\lim_{n \rightarrow \infty} \mathbb{E} \exp \left\{ it \frac{M_n - \mathbb{E}M_1}{\sqrt{n}} \right\} = \exp \left\{ -\frac{t^2}{2} \right\}.$$

We showed the pointwise convergence of characteristic functions to the characteristic function of the standard normal distribution. This proves the desired convergence in distribution. \square

At the end we present (without proof) a generalization of the Feller-Lindeberg central limit theorem for triangular array of martingale differences.

Theorem 5.4. (Brown's central limit theorem for martingale differences) *Consider a triangular array $\{D_{k,n}, k = 1, \dots, k_n, n \in \mathbb{N}\}$ such that for each row $n \in \mathbb{N}$ there are σ -algebras $\mathcal{F}_{0,n} = \{\emptyset, \Omega\} \subseteq \mathcal{F}_{1,n} \subseteq \mathcal{F}_{2,n} \subseteq \dots \subseteq \mathcal{F}_{k_n,n}$ and $(D_{1,n}, \dots, D_{k_n,n})$ is $\{\mathcal{F}_{k,n}, k = 1, \dots, k_n\}$ -martingale difference sequence. Assume that*

(i) *the conditional Feller-Lindeberg condition is satisfied, that is,*

$$\sum_{k=1}^{k_n} \mathbb{E} [D_{k,n}^2 \mathbf{1}_{\{|D_{k,n}| \geq \varepsilon\}} | \mathcal{F}_{k-1,n}] \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 0 \quad \text{for every } \varepsilon > 0,$$

(ii) $\sum_{k=1}^{k_n} \mathbb{E}[D_{k,n}^2 | \mathcal{F}_{k-1,n}] \xrightarrow[n \rightarrow \infty]{\mathbb{P}} 1$.

Then

$$\sum_{k=1}^{k_n} D_{k,n} \xrightarrow[n \rightarrow \infty]{d} N(0, 1).$$

Note that Theorem 5.3 is a special case of Theorem 5.4. It suffices to take $D_{k,n} = D_k/\sqrt{n}$, $k = 1, \dots, k_n = n$. The Conditional Feller-Lindeberg condition follows from the assumption that the conditional absolute third moments are bounded.